




ELSEVIER

Contents lists available at ScienceDirect

## Chinese Journal of Physics

journal homepage: [www.sciencedirect.com/journal/chinese-journal-of-physics](http://www.sciencedirect.com/journal/chinese-journal-of-physics)

# Solitons and conservation laws in magneto–optic waveguides with generalized Kudryashov’s equation

Elsayed M.E. Zayed<sup>a</sup>, Mohamed E.M. Alngar<sup>a</sup>, Anjan Biswas<sup>b,c,d</sup>, Abdul H. Kara<sup>e</sup>,  
Mir Asma<sup>f</sup>, Mehmet Ekici<sup>\*g</sup>, Salam Khan<sup>b</sup>, Abdullah K. Alzahrani<sup>c</sup>,  
Milivoj R. Belic<sup>h</sup>

<sup>a</sup> Mathematics Department, Faculty of Science, Zagazig University, Zagazig 44519, Egypt

<sup>b</sup> Department of Physics, Chemistry and Mathematics, Alabama A&M University, Normal, AL 35762–4900, USA

<sup>c</sup> Department of Mathematics, King Abdulaziz University, Jeddah 21589, Saudi Arabia

<sup>d</sup> Department of Applied Mathematics, National Research Nuclear University, 31 Kashirskoe Shosse, Moscow 115409, Russian Federation

<sup>e</sup> School of Mathematics, University of the Witwatersrand, Private Bag 3, Johannesburg, Wits-2050, South Africa

<sup>f</sup> Institute of Mathematical Sciences, Faculty of Science, University of Malaya, Kuala Lumpur 50603, Malaysia

<sup>g</sup> Department of Mathematics, Faculty of Science and Arts, Yozgat Bozok University, Yozgat 66100, Turkey

<sup>h</sup> Science Program, Texas A&M University at Qatar, PO Box 23874, Doha, Qatar

## ARTICLE INFO

## Keywords:

Solitons

Magneto–optic

Generalized Kudryashov’s equation

## OCIS:

060.2310

060.4510

060.5530

190.3270

190.4370

## ABSTRACT

The modified sub–ODE approach secures optical soliton solutions in magneto–optic waveguides with generalized Kudryashov’s equation. The solutions are initially drafted in terms of Jacobi’s elliptic functions. The limiting process, when the modulus of ellipticity approaches zero or unity, the soliton solutions emerge. A few solutions in terms of Weierstrass’ elliptic functions are also revealed. Finally, the conservation laws are computed for the model using the multiplier approach.

## 1. Introduction

An important application of optical solitons to a wide range of waveguides is in magneto–optic waveguides [1–20]. This carries several advantages such as controlling soliton cluster and thus transferring these soliton molecules to a state of “separation”. This addresses a growing problem in the age of telecommunications, namely Internet bottleneck. Thus, it is imperative and vital to take a deeper look at soliton dynamics in such waveguides.

During 2019, Kudryashov proposed a new form of intensity–dependent refractive index that led to a wide range of reported results [9]. Later during 2020, this law was generalized and extended with eight nonlinear terms that gave way to what is being referred to as generalized Kudryashov’s equation [10]. Today’s paper implements this generalized Kudryashov’s law of refractive index to magneto–optic waveguides and study the dynamics of solitons in it. Our approach in today’s work is the sub–ODE scheme. This algorithm enables the governing model to yield bright, dark and singular soliton solutions to the model. The existence criteria of these solitons are

\* Corresponding author.

E-mail address: [mehmet.ekici@bozok.edu.tr](mailto:mehmet.ekici@bozok.edu.tr) (M. Ekici).

<https://doi.org/10.1016/j.cjph.2020.11.026>

Received 13 July 2020; Received in revised form 1 November 2020; Accepted 7 November 2020

Available online 26 December 2020

0577-9073/© 2020 The Physical Society of the Republic of China (Taiwan). Published by Elsevier B.V. All rights reserved.

also captured from the structure of the soliton solutions. Finally, the conservation laws and the conserved quantities are also recovered for the model. This gave way to the zone of existence for the solitons that depends on the power law nonlinearity parameter in Kudryashov’s law of refractive index. The results along with the detailed analytical perspectives are all exhibited in the rest of the paper.

### 1.1. Governing model

The governing nonlinear Schrödinger’s equation (NLSE) for the generalized Kudryashov’s equation in polarization preserving fibers is written as [10]:

$$iq_t + ia_1q_x + a_2q_{xx} + ia_3q_{xxx} + a_4q_{xxxx} + \sigma q + \left( \frac{\nu_4}{|q|^{4n}} + \frac{\nu_3}{|q|^{3n}} + \frac{\nu_2}{|q|^{2n}} + \frac{\nu_1}{|q|^n} + \mu_1|q|^n + \mu_2|q|^{2n} + \mu_3|q|^{3n} + \mu_4|q|^{4n} \right) q = 0 \tag{1}$$

where  $\sigma, a_m, \nu_m, \mu_m, (m = 1, 2, 3, 4)$  are parameters, while  $i = \sqrt{-1}$ . The dependent variable  $q = q(x, t)$  is the complex-valued describing the pulse profile. The independent variables  $x$  and  $t$  represent spatial and temporal variables, respectively. The nonlinearity index  $n$  is the power parameter.

The coupled system of NLSE for the generalized Kudryashov’s equation in magneto-optic waveguides is written, for the first time, in the form:

$$\begin{aligned} &iu_t + ia_1u_x + b_1u_{xx} + ic_1u_{xxx} + d_1u_{xxxx} + \sigma_1u \\ &+ \left( \frac{n_1}{|u|^{4n}} + \frac{m_1}{|u|^{3n}} + \frac{l_1}{|u|^{2n}} + \frac{k_1}{|u|^n} + e_1|u|^n + f_1|u|^{2n} + g_1|u|^{3n} + h_1|u|^{4n} \right) u \\ &+ \left( \frac{\tau_1}{|v|^{4n}} + \frac{\zeta_1}{|v|^{3n}} + \frac{\eta_1}{|v|^{2n}} + \frac{\xi_1}{|v|^n} + \alpha_1|v|^n + \beta_1|v|^{2n} + \gamma_1|v|^{3n} + \delta_1|v|^{4n} \right) u \\ &= Q_1v + i \left[ \lambda_1 \left( |u|^{2n}u \right)_x + \nu_1 \left( |u|^{2n} \right)_x u + \theta_1|u|^{2n}u_x \right] \\ &iv_t + ia_2v_x + b_2v_{xx} + ic_2v_{xxx} + d_2v_{xxxx} + \sigma_2v \\ &+ \left( \frac{n_2}{|v|^{4n}} + \frac{m_2}{|v|^{3n}} + \frac{l_2}{|v|^{2n}} + \frac{k_2}{|v|^n} + e_2|v|^n + f_2|v|^{2n} + g_2|v|^{3n} + h_2|v|^{4n} \right) v \\ &+ \left( \frac{\tau_2}{|u|^{4n}} + \frac{\zeta_2}{|u|^{3n}} + \frac{\eta_2}{|u|^{2n}} + \frac{\xi_2}{|u|^n} + \alpha_2|u|^n + \beta_2|u|^{2n} + \gamma_2|u|^{3n} + \delta_2|u|^{4n} \right) v \\ &= Q_2u + i \left[ \lambda_2 \left( |v|^{2n}v \right)_x + \nu_2 \left( |v|^{2n} \right)_x v + \theta_2|v|^{2n}v_x \right] \end{aligned} \tag{2}$$

where  $a_j, b_j, c_j, d_j, \sigma_j, e_j, f_j, g_j, h_j, k_j, l_j, m_j, n_j, \alpha_j, \beta_j, \gamma_j, \delta_j, \xi_j, \eta_j, \zeta_j, \tau_j, Q_j, \lambda_j, \nu_j$  and  $\theta_j$  for  $j = 1, 2$  are all real constants, while  $i = \sqrt{-1}$ . The system (2) and (3) are coupled NLSEs for the generalization of Kudryashov’s equation that govern the propagation of optical solitons through magneto-optic waveguides, where the independent variables  $x$  and  $t$  represent the spatial and temporal variables, respectively. The dependent variables  $u(x, t)$  and  $v(x, t)$  are the complex valued wave profiles, where  $a_j$  represent the coefficients of inter-modal dispersion (IMD),  $b_j$  are the coefficients of group velocity dispersion (GVD),  $c_j$  are the coefficients of third-order dispersion (3OD) and  $d_j$  are the coefficients of fourth-order dispersion (4OD). Next,  $e_j, f_j, g_j, h_j, k_j, l_j, m_j$  and  $n_j$  are the coefficients of self-phase modulation (SPM). The parameters  $\alpha_j, \beta_j, \gamma_j, \delta_j, \xi_j, \eta_j, \zeta_j$  and  $\tau_j$  represent the cross-phase modulation (XPM). Here  $\sigma_j$  come from the detuning effect and  $Q_j$  are the coefficients of magneto-optic parameters. The parameters  $\lambda_j$  give the self-steepening (SS) terms that avoid the formation of shock waves. Finally  $\nu_j$  and  $\theta_j$  stand for the coefficients of nonlinear dispersion.

The objective of today’s work is to reveal bright, dark and singular soliton solutions in magneto-optic waveguides with generalized Kudryashov’s equation by the sub-ODE scheme and then compute the conservation laws for the model employing the multiplier approach. It worths noting that the results recovered are new and have not been reported before.

This article is organized as follows: In Section 2, mathematical analysis is discussed. In Section 3 we solve Eqs. (2) and (3) using the modified sub-ODE method. In Section 4, the conservation laws are displayed while finally in Section 5, conclusions are drawn.

## 2. Mathematical analysis

In order to solve Eqs. (2) and (3), we assume that the hypothesis as:

$$u(x, t) = \phi_1(\xi)\exp[i\psi(x, t)] \tag{4}$$

$$v(x, t) = \phi_2(\xi)\exp[i\psi(x, t)] \tag{5}$$

and

$$\xi = x - ct, \quad \psi(x, t) = -\kappa x + \omega t + \theta_0 \tag{6}$$

where  $c, \kappa, \omega$  and  $\theta_0$  are all non zero constants to be determined which represent velocity of soliton, frequency of soliton, wave number and phase constant, respectively, while  $\phi_1(\xi), \phi_2(\xi)$  and  $\psi(x, t)$  are real functions which represent the amplitude portion of the soliton and the phase component of the soliton, respectively. Substituting (4) and (5) along with (6) into Eqs. (2) and (3), separating the real and imaginary parts, we deduce that the real parts are

$$d_1\phi_1'''' + (b_1 + 3c_1\kappa - 6d_1\kappa^2)\phi_1'' + (\sigma_1 - \omega + a_1\kappa - b_1\kappa^2 - c_1\kappa^3 + d_1\kappa^4)\phi_1 - Q_1\phi_2 - \kappa(\lambda_1 + \theta_1)\phi_1^{2n+1} + e_1\phi_1^{1+n} + f_1\phi_1^{1+2n} + g_1\phi_1^{1+3n} + h_1\phi_1^{1+4n} + k_1\phi_1^{1-n} + l_1\phi_1^{1-2n} + m_1\phi_1^{1-3n} + n_1\phi_1^{1-4n} + (\alpha_1\phi_2^n + \beta_1\phi_2^{2n} + \gamma_1\phi_2^{3n} + \delta_1\phi_2^{4n} + \xi_1\phi_2^{-n} + \eta_1\phi_2^{-2n} + \zeta_1\phi_2^{-3n} + \tau_1\phi_2^{-4n})\phi_1 = 0 \tag{7}$$

$$d_2\phi_2'''' + (b_2 + 3c_2\kappa - 6d_2\kappa^2)\phi_2'' + (\sigma_2 - \omega + a_2\kappa - b_2\kappa^2 - c_2\kappa^3 + d_2\kappa^4)\phi_2 - Q_2\phi_1 - \kappa(\lambda_2 + \theta_2)\phi_2^{2n+1} + e_2\phi_2^{1+n} + f_2\phi_2^{1+2n} + g_2\phi_2^{1+3n} + h_2\phi_2^{1+4n} + k_2\phi_2^{1-n} + l_2\phi_2^{1-2n} + m_2\phi_2^{1-3n} + n_2\phi_2^{1-4n} + (\alpha_2\phi_1^n + \beta_2\phi_1^{2n} + \gamma_2\phi_1^{3n} + \delta_2\phi_1^{4n} + \xi_2\phi_1^{-n} + \eta_2\phi_1^{-2n} + \zeta_2\phi_1^{-3n} + \tau_2\phi_1^{-4n})\phi_2 = 0 \tag{8}$$

while the imaginary parts are

$$(c_1 - 4d_1\kappa)\phi_1'' + (a_1 - c - 2b_1\kappa - 3c_1\kappa^2 + 4d_1\kappa^3)\phi_1' - [(2n + 1)\lambda_1 + 2n\nu_1 + \theta_1]\phi_1^{2n}\phi_1' = 0 \tag{9}$$

$$(c_2 - 4d_2\kappa)\phi_2'' + (a_2 - c - 2b_2\kappa - 3c_2\kappa^2 + 4d_2\kappa^3)\phi_2' - [(2n + 1)\lambda_2 + 2n\nu_2 + \theta_2]\phi_2^{2n}\phi_2' = 0. \tag{10}$$

The linearly independent principle is applied on (9) and (10) to get:

$$c_1 - 4d_1\kappa = 0 \tag{11}$$

$$c = a_1 - 2b_1\kappa - 8d_1\kappa^3 \tag{12}$$

$$(2n + 1)\lambda_1 + 2n\nu_1 + \theta_1 = 0 \tag{13}$$

and

$$c_2 - 4d_2\kappa = 0 \tag{14}$$

$$c = a_2 - 2b_2\kappa - 8d_2\kappa^3 \tag{15}$$

$$(2n + 1)\lambda_2 + 2n\nu_2 + \theta_2 = 0. \tag{16}$$

From (12) and (15), one gets the condition:

$$a_1 - a_2 - 2\kappa(b_1 - b_2) - 8\kappa^3(d_1 - d_2) = 0. \tag{17}$$

From (11) and (14), we deduce that the frequency of soliton  $\kappa$  is given by:

$$\kappa = \frac{c_j}{4d_j} \tag{18}$$

provided  $d_j \neq 0$  for  $j = 1, 2$ . From (18) we have the relation

$$c_1d_2 = c_2d_1. \tag{19}$$

Set

$$\phi_2(\xi) = \chi\phi_1(\xi) \tag{20}$$

where  $\chi$  is a nonzero constant, such that  $\chi \neq 1$ . Consequently, Eqs. (7) and (8) reduce to

$$\begin{aligned}
 & d_1\phi_1'''' + (b_1 + 3c_1\kappa - 6d_1\kappa^2)\phi_1'' + (\sigma_1 - \omega + a_1\kappa - b_1\kappa^2 - c_1\kappa^3 + d_1\kappa^4 - Q_1\chi)\phi_1 \\
 & + (e_1 + \alpha_1\chi^n)\phi_1^{1+n} + [f_1 + \beta_1\chi^{2n} - \kappa(\lambda_1 + \theta_1)]\phi_1^{1+2n} + (g_1 + \gamma_1\chi^{3n})\phi_1^{1+3n} \\
 & + (h_1 + \delta_1\chi^{4n})\phi_1^{1+4n} + (k_1 + \xi_1\chi^{-n})\phi_1^{1-n} + (l_1 + \eta_1\chi^{-2n})\phi_1^{1-2n} \\
 & + (m_1 + \zeta_1\chi^{-3n})\phi_1^{1-3n} + (n_1 + \tau_1\chi^{-4n})\phi_1^{1-4n} = 0
 \end{aligned} \tag{21}$$

and

$$\begin{aligned}
 & d_2\chi\phi_1'''' + (b_2 + 3c_2\kappa - 6d_2\kappa^2)\chi\phi_1'' + [(\sigma_2 - \omega + a_2\kappa - b_2\kappa^2 - c_2\kappa^3 + d_2\kappa^4)\chi - Q_2]\phi_1 \\
 & + (e_2\chi^{1+n} + \alpha_2\chi)\phi_1^{1+n} + [f_2\chi^{1+2n} + \beta_2\chi - \kappa(\lambda_2 + \theta_2)\chi^{2n+1}]\phi_1^{2n+1} + (g_2\chi^{1+3n} + \gamma_2\chi)\phi_1^{1+3n} \\
 & + (h_2\chi^{1+4n} + \delta_2\chi)\phi_1^{1+4n} + (k_2\chi^{1-n} + \xi_2\chi)\phi_1^{1-n} + (l_2\chi^{1-2n} + \eta_1\chi)\phi_1^{1-2n} \\
 & + (m_2\chi^{1-3n} + \zeta_2\chi)\phi_1^{1-3n} + (n_2\chi^{1-4n} + \tau_2\chi)\phi_1^{1-4n} = 0.
 \end{aligned} \tag{22}$$

Eqs. (21) and (2) have the same form under the constraint conditions:

$$d_1 = d_2\chi \tag{23}$$

$$b_1 + 3c_1\kappa - 6d_1\kappa^2 = (b_2 + 3c_2\kappa - 6d_2\kappa^2)\chi \tag{24}$$

$$\sigma_1 - \omega + a_1\kappa - b_1\kappa^2 - c_1\kappa^3 + d_1\kappa^4 - Q_1\chi = (\sigma_2 - \omega + a_2\kappa - b_2\kappa^2 - c_2\kappa^3 + d_2\kappa^4)\chi - Q_2 \tag{25}$$

$$e_1 + \alpha_1\chi^n = e_2\chi^{1+n} + \alpha_2\chi \tag{26}$$

$$f_1 + \beta_1\chi^{2n} - \kappa(\lambda_1 + \theta_1) = f_2\chi^{1+2n} + \beta_2\chi - \kappa(\lambda_2 + \theta_2)\chi^{2n+1} \tag{27}$$

$$g_1 + \gamma_1\chi^{3n} = g_2\chi^{1+3n} + \gamma_2\chi \tag{28}$$

$$h_1 + \delta_1\chi^{4n} = h_2\chi^{1+4n} + \delta_2\chi \tag{29}$$

$$k_1 + \xi_1\chi^{-n} = k_2\chi^{1-n} + \xi_2\chi \tag{30}$$

$$l_1 + \eta_1\chi^{-2n} = l_2\chi^{1-2n} + \eta_1\chi \tag{31}$$

$$m_1 + \zeta_1\chi^{-3n} = m_2\chi^{1-3n} + \zeta_2\chi \tag{32}$$

$$n_1 + \tau_1\chi^{-4n} = n_2\chi^{1-4n} + \tau_2\chi. \tag{33}$$

From (19), (23) and (25), the wave number  $\omega$  is given by

$$\omega = \frac{Q_2 - Q_1\chi - (\chi a_2 - a_1)\kappa + (\chi b_2 - b_1)\kappa^2 - \chi\sigma_2 + \sigma_1}{(1 - \chi)}. \tag{34}$$

Eq. (21) can be rewritten in the form:

$$\begin{aligned}
 & \phi_1'''' + H_0\phi_1'' + H_1\phi_1 + \Delta_1\phi_1^{1+n} + \Delta_2\phi_1^{1+2n} + \Delta_3\phi_1^{1+3n} + \Delta_4\phi_1^{1+4n} \\
 & + \Gamma_1\phi_1^{1-n} + \Gamma_2\phi_1^{1-2n} + \Gamma_3\phi_1^{1-3n} + \Gamma_4\phi_1^{1-4n} = 0
 \end{aligned} \tag{35}$$

where

$$\left. \begin{aligned}
 H_0 &= \frac{b_1 + 3c_1\kappa - 6d_1\kappa^2}{d_1}, & H_1 &= \frac{\sigma_1 - \omega + a_1\kappa - b_1\kappa^2 - c_1\kappa^3 + d_1\kappa^4 - Q_1\chi}{d_1}, \\
 \Delta_1 &= \frac{e_1 + \alpha_1\chi^n}{d_1}, & \Delta_2 &= \frac{f_1 + \beta_1\chi^{2n} - \kappa(\lambda_1 + \theta_1)}{d_1}, & \Delta_3 &= \frac{g_1 + \gamma_1\chi^{3n}}{d_1}, & \Delta_4 &= \frac{h_1 + \delta_1\chi^{4n}}{d_1}, \\
 \Gamma_1 &= \frac{k_1 + \xi_1\chi^{-n}}{d_1}, & \Gamma_2 &= \frac{l_1 + \eta_1\chi^{-2n}}{d_1}, & \Gamma_3 &= \frac{m_1 + \zeta_1\chi^{-3n}}{d_1}, & \Gamma_4 &= \frac{n_1 + \tau_1\chi^{-4n}}{d_1}.
 \end{aligned} \right\} \tag{36}$$

Now, the problem of finding exact solutions for Eqs. (2) and (3) is transformed for solving Eq. (2).

### 3. The modified sub-ode method

Let us demonstrate that there is a solution of Eq. (2) in the form

$$\phi_1(\xi) = F(\xi) \tag{37}$$

where  $F(\xi)$  is the solution of the following modified sub-ODE equation [20–25]:

$$F'^2(\xi) = AF^{2-2n}(\xi) + BF^{2-n}(\xi) + CF^2(\xi) + DF^{2+n}(\xi) + EF^{2+2n}(\xi), \quad n > 0 \tag{38}$$

where  $A, B, C, D$  and  $E$  are constants. We will use (37) and (38) to get

$$\phi_1''(\xi) = (1-n)AF^{1-2n}(\xi) + \frac{1}{2}(2-n)BF^{1-n}(\xi) + CF(\xi) + \frac{1}{2}(2+n)DF^{1+n}(\xi) + (1+n)EF^{1+2n}(\xi) \tag{39}$$

$$\begin{aligned} \phi_1''(\xi) = & \left[ (1-n)(1-2n)AF^{-2n}(\xi) + \frac{1}{2}(2-n)(1-n)BF^{-n}(\xi) + C \right. \\ & \left. + \frac{1}{2}(2+n)(1+n)DF^n(\xi) + (1+n)(1+2n)EF^{2n}(\xi) \right] F_1'(\xi) \end{aligned} \tag{40}$$

and

$$\begin{aligned} \phi_1'''(\xi) = & (1-n) \left[ \frac{n^2}{4}(3B^2 + 16AC) + (1-2n)(2AC + B^2) \right] F^{1-2n}(\xi) \\ & + (2-n) \left[ \frac{n^2}{2}(BC + 6AD) + (1-n)(BC + AD) \right] F^{1-n}(\xi) \\ & + (1+n) \left[ \frac{n^2}{4}(3D^2 + 16EC) + (2n+1)(D^2 + 2EC) \right] F^{1+2n}(\xi) \\ & + (3n-1)(2n-1)(1-n)A^2F^{1-4n}(\xi) + (3n-2)(2n-1)(1-n)ABF^{1-3n}(\xi) \\ & + (3n+2)(2n+1)(1+n)EDF^{1+3n}(\xi) + (2n+1)(3n+1)(1+n)E^2F^{1+4n}(\xi) \\ & + (2+n) \left[ \frac{n^2}{2}(DC + 6BE) + (DC + BE)(n+1) \right] F^{1+n}(\xi) \\ & + \left[ \frac{5n^2}{2}(4AE + DB) + 2(AE + DB) + C^2 \right] F(\xi). \end{aligned} \tag{41}$$

Substituting (37), (39) and (41) into Eq. (2), collecting all the coefficients of  $[F(\xi)]^{Jn+1}[F'(\xi)]^s$ , ( $J = 0, 1, 2, \dots, 8, s = 0, 1$ ) and setting them to zero, one gets the following algebraic equations:

$$\left. \begin{aligned} (3n+1)(2n+1)(1+n)E^2 + \Delta_4 &= 0, \\ (3n+2)(2n+1)(1+n)ED + \Delta_3 &= 0, \\ (1+n) \left[ n^2 \left( \frac{3}{4}D^2 + 4EC \right) + (2n+1)(D^2 + 2EC) \right] + (1+n)H_0E + \Delta_2 &= 0, \\ (2+n) \left[ (6BE + DC)n^2 + 2(n+1)(BE + DC) \right] + (n+2)DH_0 + 2\Delta_1 &= 0, \\ \frac{5n^2}{2}(4AE + DB) + 2(DB + AE) + C^2 + H_0C + H_1 &= 0, \\ (n-2) \left[ (6AD + BC)n^2 + 2(1-n)(BC + AD) \right] + (n-2)H_0B - 2\Gamma_1 &= 0, \\ (1-n) \left[ \left( \frac{3}{4}B^2 + 4AC \right) n^2 + (1-2n)(2AC + B^2) \right] + \Gamma_2 + (1-n)AH_0 &= 0, \\ (2n-1)(3n-2)(1-n)BA + \Gamma_3 &= 0, \\ (2n-1)(3n-1)(1-n)A^2 + \Gamma_4 &= 0. \end{aligned} \right\} \tag{42}$$

Solving the above algebraic Eq. (42), one gets the following conditions for existence of the solutions  $\phi_1(\xi) = F(\xi)$  of Eq. (2):

$$\Delta_1 = -(2+n) \left[ \left( 3BE + \frac{1}{2}DC \right) n^2 + (n+1)(BE + DC) + \frac{1}{2}DH_0 \right] \tag{43}$$

$$\Delta_2 = -(1+n) \left[ n^2 \left( \frac{3}{4}D^2 + 4EC \right) + (2n+1)(D^2 + 2EC) + H_0E \right] \tag{44}$$

$$\Delta_3 = -(3n+2)(2n+1)(1+n)ED \tag{45}$$

$$\Delta_4 = -(3n + 1)(2n + 1)(1 + n)E^2 \tag{46}$$

$$H_1 = -5n^2 \left( 2AE + \frac{1}{2}DB \right) - 2(DB + AE) - C^2 - H_0C \tag{47}$$

$$\Gamma_1 = \frac{(n - 2)}{2} [(6AD + BC)n^2 + 2(1 - n)(BC + AD) + H_0B] \tag{48}$$

$$\Gamma_2 = (n - 1) \left[ \left( \frac{3}{4}B^2 + 4AC \right) n^2 + (1 - 2n)(2AC + B^2) + AH_0 \right] \tag{49}$$

$$\Gamma_3 = (2n - 1)(3n - 2)(n - 1)BA \tag{50}$$

$$\Gamma_4 = (2n - 1)(3n - 1)(n - 1)A^2. \tag{51}$$

According to the modified Sub-ODE method, we have the following types of solutions for Eqs. (2) and (3):

**Case-1:** If we substitute  $A = B = D = 0$  in the above conditions (43)–(51), then one gets the results:

$$H_1 = -C(C + H_0), \quad \Delta_1 = 0, \quad \Delta_2 = -(1 + n)E[(4n^2 + 4n + 2)C + H_0], \quad \Delta_3 = 0, \\ \Delta_4 = -(3n + 1)(2n + 1)(1 + n)E^2, \quad \Gamma_1 = 0, \quad \Gamma_2 = 0, \quad \Gamma_3 = 0, \quad \Gamma_4 = 0. \tag{52}$$

Under the constraint conditions (52), we have the solutions:

(I) The bright soliton solutions:

$$u(x, t) = \left[ \varepsilon \sqrt{\frac{-C}{E}} \operatorname{sech}(n\sqrt{C}(x - ct)) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{53}$$

$$v(x, t) = \chi \left[ \varepsilon \sqrt{\frac{-C}{E}} \operatorname{sech}(n\sqrt{C}(x - ct)) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{54}$$

provided  $\varepsilon = \pm 1, C > 0$  and  $E < 0$ .

(II) The periodic solutions:

$$u(x, t) = \left[ \varepsilon \sqrt{\frac{C}{E}} \sec(n\sqrt{-C}(x - ct)) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{55}$$

$$v(x, t) = \chi \left[ \varepsilon \sqrt{\frac{C}{E}} \sec(n\sqrt{-C}(x - ct)) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{56}$$

provided  $\varepsilon = \pm 1, C < 0$  and  $E > 0$ .

(III) The rational solutions:

$$u(x, t) = \left[ \frac{\varepsilon}{n\sqrt{E}(x - ct)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{57}$$

$$v(x, t) = \chi \left[ \frac{\varepsilon}{n\sqrt{E}(x - ct)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{58}$$

provided  $\varepsilon = \pm 1, C = 0$  and  $E > 0$ .

**Case-2:** If we substitute  $B = D = 0, A = \frac{C^2}{4E}$ , in the above conditions (43)–(51), then one gets the results:

$$H_1 = -\left[ \frac{1}{2}(5n^2 + 3)C + H_0 \right] C, \quad \Delta_1 = 0, \quad \Delta_2 = -(1 + n)E[(4n^2 + 4n + 2)C + H_0], \quad \Delta_3 = 0, \\ \Delta_4 = -(3n + 1)(2n + 1)(1 + n)E^2, \quad \Gamma_1 = 0, \quad \Gamma_2 = \frac{(n - 1)C^2}{4E} [C(4n^2 - 4n + 2) + H_0], \\ \Gamma_3 = 0, \quad \Gamma_4 = \frac{C^4}{16E^2} (2n - 1)(3n - 1)(n - 1). \tag{59}$$

Under the constraint conditions (59), we have the solutions:

(I) The dark soliton solutions:

$$u(x, t) = \left[ \varepsilon \sqrt{\frac{C}{2E}} \tanh \left( n \sqrt{\frac{C}{2}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{60}$$

$$v(x, t) = \chi \left[ \varepsilon \sqrt{\frac{C}{2E}} \tanh \left( n \sqrt{\frac{C}{2}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{61}$$

provided  $\varepsilon = \pm 1, C < 0$  and  $E > 0$ .

**(II) The periodic solutions:**

$$u(x, t) = \left[ \varepsilon \sqrt{\frac{C}{2E}} \tan \left( n \sqrt{\frac{C}{2}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{62}$$

$$v(x, t) = \chi \left[ \varepsilon \sqrt{\frac{C}{2E}} \tan \left( n \sqrt{\frac{C}{2}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{63}$$

provided  $\varepsilon = \pm 1, C > 0$  and  $E > 0$ .

**Case-3:** If we substitute  $B = D = 0$ , in the above conditions (43)–(51), then, one arrives at the results:

$$\begin{aligned} H_1 &= -[2(5n^2 + 1)EA + (C + H_0)C], & \Delta_1 &= 0, & \Delta_2 &= -(1 + n)E[2C(2n^2 + 2n + 1) + H_0], \\ \Delta_3 &= 0, & \Delta_4 &= -(3n + 1)(2n + 1)(1 + n)E^2, & \Gamma_2 &= A(n - 1)[2C(2n^2 - 2n + 1) + H_0], \\ \Gamma_1 &= 0, & \Gamma_3 &= 0, & \Gamma_4 &= (2n - 1)(3n - 1)(n - 1)A^2. \end{aligned} \tag{64}$$

Under the constraint conditions (64), we have the solutions:

**(I) If  $A = \frac{m^2(m^2 - 1)C^2}{E(2m^2 - 1)^2}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:**

$$u(x, t) = \left[ \varepsilon \sqrt{\frac{Cm^2}{E(2m^2 - 1)}} \operatorname{cn} \left( n \sqrt{\frac{C}{2m^2 - 1}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{65}$$

$$v(x, t) = \chi \left[ \varepsilon \sqrt{\frac{Cm^2}{E(2m^2 - 1)}} \operatorname{cn} \left( n \sqrt{\frac{C}{2m^2 - 1}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{66}$$

provided

$$\varepsilon = \pm 1, C(2m^2 - 1) > 0 \text{ and } E < 0. \tag{67}$$

**(II) If  $A = \frac{C^2(1 - m^2)}{E(2 - m^2)^2}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:**

$$u(x, t) = \left[ \varepsilon \sqrt{\frac{C}{E(2 - m^2)}} \operatorname{dn} \left( n \sqrt{\frac{C}{2 - m^2}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{68}$$

$$v(x, t) = \chi \left[ \varepsilon \sqrt{\frac{C}{E(2 - m^2)}} \operatorname{dn} \left( n \sqrt{\frac{C}{2 - m^2}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{69}$$

provided  $\varepsilon = \pm 1, C > 0$  and  $E < 0$ .

**(III) If  $A = \frac{C^2 m^2}{E(m^2 + 1)^2}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:**

$$u(x, t) = \left[ \varepsilon \sqrt{\frac{Cm^2}{E(m^2 + 1)}} \operatorname{sn} \left( n \sqrt{\frac{C}{m^2 + 1}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{70}$$

$$v(x, t) = \chi \left[ \varepsilon \sqrt{\frac{Cm^2}{E(m^2 + 1)}} \operatorname{sn} \left( n \sqrt{\frac{C}{m^2 + 1}} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{71}$$

provided  $\varepsilon = \pm 1, C < 0$  and  $E > 0$ .

**(IV) The Weierstrass elliptic function solutions**

$$u(x, t) = \left[ \frac{1}{E} \wp(n(x - ct), g_2, g_3) - \frac{C}{3E} \right]^{\frac{1}{2n}} e^{i(-kx + \omega t + \theta_0)} \tag{72}$$

$$v(x, t) = \chi \left[ \frac{1}{E} \wp(n(x - ct), g_2, g_3) - \frac{C}{3E} \right]^{\frac{1}{2n}} e^{i(-kx + \omega t + \theta_0)} \tag{73}$$

or

$$u(x, t) = \left[ \frac{3A}{3\wp(n(x - ct), g_2, g_3) - C} \right]^{\frac{1}{2n}} e^{i(-kx + \omega t + \theta_0)} \tag{74}$$

$$v(x, t) = \chi \left[ \frac{3A}{3\wp(n(x - ct), g_2, g_3) - C} \right]^{\frac{1}{2n}} e^{i(-kx + \omega t + \theta_0)} \tag{75}$$

where

$$g_2 = \frac{4C^2 - 12AE}{3} \quad \text{and} \quad g_3 = \frac{4C(-2C^2 + 9AE)}{27}. \tag{76}$$

Here  $\wp(\xi, g_2, g_3)$  is called a Weierstrass elliptic function which satisfies the Eq.  $\wp'^2 = 4\wp^3 - g_2\wp - g_3$ , such that  $g_2$  and  $g_3$  are called invariants of the Weierstrass elliptic function, in which  $\xi = \frac{d}{d\xi}$ .

(V) The Weierstrass elliptic function solutions

$$u(x, t) = \left[ \frac{6\sqrt{A}\wp(n(x - ct), g_2, g_3) + C\sqrt{A}}{3\wp'(n(x - ct), g_2, g_3)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{77}$$

$$v(x, t) = \chi \left[ \frac{6\sqrt{A}\wp(n(x - ct), g_2, g_3) + C\sqrt{A}}{3\wp'(n(x - ct), g_2, g_3)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{78}$$

or

$$u(x, t) = \left[ \frac{3\sqrt{E^{-1}}\wp'(n(x - ct), g_2, g_3)}{6\wp(n(x - ct), g_2, g_3) + C} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{79}$$

$$v(x, t) = \chi \left[ \frac{3\sqrt{E^{-1}}\wp'(n(x - ct), g_2, g_3)}{6\wp(n(x - ct), g_2, g_3) + C} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{80}$$

provided  $A > 0$  and  $E > 0$ , where

$$g_2 = \frac{C^2}{12} + AE \quad \text{and} \quad g_3 = \frac{C(36AE - C^2)}{216}. \tag{81}$$

**Case-4:** If we substitute  $A = B = E = 0$ , in the above conditions (43)–(51), then, one gets the results:

$$\begin{aligned} H_1 &= -C(C + H_0), \quad \Delta_1 = -\frac{(2+n)D}{2} [C(n^2 + 2n + 2) + H_0], \\ \Delta_2 &= -\frac{1}{4}(1+n)(3n^2 + 8n + 4)D^2, \quad \Delta_3 = 0, \quad \Delta_4 = 0, \\ \Gamma_1 &= 0, \quad \Gamma_2 = 0, \quad \Gamma_3 = 0, \quad \Gamma_4 = 0. \end{aligned} \tag{82}$$

Under the constraint conditions (82), we have the solutions:

(I) The bright soliton solutions:

$$u(x, t) = \left[ -\frac{C}{D} \operatorname{sech}^2\left(\frac{n}{2}\sqrt{C}(x - ct)\right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{83}$$

$$v(x, t) = \chi \left[ -\frac{C}{D} \operatorname{sech}^2\left(\frac{n}{2}\sqrt{C}(x - ct)\right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{84}$$

provided  $C > 0$  and  $D < 0$ .

(II) The periodic solutions:



$$u(x, t) = \left[ -\frac{C}{D} \operatorname{sec}^2 \left( \frac{n}{2} \sqrt{-C} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{85}$$

$$v(x, t) = \chi \left[ -\frac{C}{D} \operatorname{sec}^2 \left( \frac{n}{2} \sqrt{-C} (x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{86}$$

provided  $C < 0$  and  $D > 0$ .

**(III)** The rational solutions:

$$u(x, t) = \left[ \frac{4}{n^2 D (x - ct)^2} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{87}$$

$$v(x, t) = \chi \left[ \frac{4}{n^2 D (x - ct)^2} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{88}$$

provided  $C = 0$  and  $D > 0$ . **Case-5:** If we substitute  $C = E = 0$ , in the above conditions (43)–(51), then, one gets the results:

$$\begin{aligned} H_1 &= -\frac{1}{2}(5n^2 + 4)DB, & \Delta_1 &= -\frac{1}{2}(2 + n)DH_0, \\ \Delta_2 &= -\frac{1}{4}D^2(1 + n)(3n^2 + 8n + 4), & \Delta_3 &= 0, & \Delta_4 &= 0, \\ \Gamma_1 &= \frac{(n - 2)}{2} [(6n^2 - 2n + 2)AD + H_0B], & \Gamma_2 &= \frac{(n - 1)}{4} [B^2(3n - 2)(n - 2) + 4AH_0], \\ \Gamma_3 &= (2n - 1)(3n - 2)(n - 1)BA, & \Gamma_4 &= (2n - 1)(3n - 1)(n - 1)A^2. \end{aligned} \tag{89}$$

Under the constraint conditions (89), we have the solutions:

$$u(x, t) = \left[ \wp \left( \frac{n}{2} \sqrt{D} (x - ct), -\frac{4B}{D}, -\frac{4A}{D} \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{90}$$

$$v(x, t) = \chi \left[ \wp \left( \frac{n}{2} \sqrt{D} (x - ct), -\frac{4B}{D}, -\frac{4A}{D} \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{91}$$

provided  $D > 0$ .

**Case-6:** If we substitute  $A = B = 0$  in the above conditions (43)–(51), then one gets the results:

$$\begin{aligned} H_1 &= -C(C + H_0), & \Delta_1 &= -\frac{(2 + n)}{2} D [C(n^2 + 2n + 2) + H_0], \\ \Delta_2 &= -(1 + n) \left\{ \left[ n^2 \left( \frac{3}{4} D^2 + 4EC \right) + (2n + 1)(D^2 + 2EC) \right] + H_0 E \right\}, \\ \Delta_3 &= -(3n + 2)(2n + 1)(1 + n)ED, & \Delta_4 &= -(3n + 1)(2n + 1)(1 + n)E^2, \\ \Gamma_1 &= 0, & \Gamma_2 &= 0, & \Gamma_3 &= 0, & \Gamma_4 &= 0. \end{aligned} \tag{92}$$

Under the constraint conditions (92), we have the solutions:

**(I)** If  $E = \frac{D^2}{4C} - C$  and  $C > 0$ , then one gets the bright soliton solutions:

$$u(x, t) = \left[ \frac{1}{\cosh(n\sqrt{C}(x - ct)) - \frac{D}{2C}} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{93}$$

$$v(x, t) = \chi \left[ \frac{1}{\cosh(n\sqrt{C}(x - ct)) - \frac{D}{2C}} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{94}$$

provided  $D < 2C$ .

**(II)** If  $D = -2\sqrt{CE}$  and  $C > 0$ , then one gets the dark soliton solutions:

$$u(x, t) = \left[ \frac{1}{2} \sqrt{\frac{C}{E}} \left[ 1 + \epsilon \tanh \left( \frac{n}{2} \sqrt{C} (x - ct) \right) \right] \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{95}$$

$$v(x, t) = \chi \left[ \frac{1}{2} \sqrt{\frac{C}{E}} \left[ 1 + \varepsilon \tanh\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right] \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{96}$$

provided  $\varepsilon = \pm 1$  and  $E > 0$ .

(III) If  $C = 0$ , then one gets the rational solutions:

$$u(x, t) = \left[ \frac{4D}{[nD(x - ct)]^2 - 4E} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{97}$$

$$v(x, t) = \chi \left[ \frac{4D}{[nD(x - ct)]^2 - 4E} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{98}$$

provided  $E < 0$  and  $D > 0$ .

(IV) If  $D^2 - 4CE > 0$  and  $C > 0$ , then one gets the bright soliton solutions:

$$u(x, t) = \left[ \frac{2C \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{2\sqrt{D^2 - 4CE} - [\sqrt{D^2 - 4CE} + D] \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{99}$$

$$v(x, t) = \chi \left[ \frac{2C \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{2\sqrt{D^2 - 4CE} - [\sqrt{D^2 - 4CE} + D] \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{100}$$

or

$$u(x, t) = \left[ \frac{2C}{\varepsilon \sqrt{D^2 - 4CE} \cosh(n\sqrt{C}(x - ct)) - D} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{101}$$

$$v(x, t) = \chi \left[ \frac{2C}{\varepsilon \sqrt{D^2 - 4CE} \cosh(n\sqrt{C}(x - ct)) - D} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{102}$$

and the singular soliton solutions:

$$u(x, t) = \left[ \frac{2C \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{2\sqrt{D^2 - 4CE} + [\sqrt{D^2 - 4CE} - D] \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{103}$$

$$v(x, t) = \chi \left[ \frac{2C \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{2\sqrt{D^2 - 4CE} + [\sqrt{D^2 - 4CE} - D] \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{104}$$

provided  $\varepsilon = \pm 1$ .

(V) If  $D^2 - 4CE < 0$  and  $C > 0$ , then one gets the singular soliton solutions:

$$u(x, t) = \left[ \frac{2C}{\varepsilon \sqrt{-(D^2 - 4CE)} \sinh(n\sqrt{C}(x - ct)) - D} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{105}$$

$$v(x, t) = \chi \left[ \frac{2C}{\varepsilon \sqrt{-(D^2 - 4CE)} \sinh(n\sqrt{C}(x - ct)) - D} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{106}$$

provided  $\varepsilon = \pm 1$ .

(VI) If  $D^2 - 4CE = 0$  and  $C > 0$ , then one gets the bright soliton solutions:

$$u(x, t) = \left[ -\frac{C}{D} \left\{ 1 + \varepsilon \tanh\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{107}$$

$$v(x, t) = \chi \left[ -\frac{C}{D} \left\{ 1 + \varepsilon \tanh\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{108}$$

and the singular soliton solutions:

$$u(x, t) = \left[ -\frac{C}{D} \left\{ 1 + \varepsilon \coth\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{109}$$

$$v(x, t) = \chi \left[ -\frac{C}{D} \left\{ 1 + \varepsilon \coth\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{110}$$

provided  $\varepsilon = \pm 1$ .

**(VII)** If  $C > 0$ , one gets the solitary wave solutions:

$$u(x, t) = \left[ -\frac{CD \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D^2 - CE \left[ 1 + \varepsilon \tanh\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right]^2} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{111}$$

$$v(x, t) = \chi \left[ -\frac{CD \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D^2 - CE \left[ 1 + \varepsilon \tanh\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right]^2} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{112}$$

or

$$u(x, t) = \left[ \frac{CD \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D^2 - CE \left[ 1 + \varepsilon \coth\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right]^2} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{113}$$

$$v(x, t) = \chi \left[ \frac{CD \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D^2 - CE \left[ 1 + \varepsilon \coth\left(\frac{n}{2} \sqrt{C}(x - ct)\right) \right]^2} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{114}$$

or

$$u(x, t) = \left[ -\frac{C \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D + 2\varepsilon \sqrt{CE} \tanh\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{115}$$

$$v(x, t) = \chi \left[ -\frac{C \operatorname{sech}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D + 2\varepsilon \sqrt{CE} \tanh\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{116}$$

or

$$u(x, t) = \left[ \frac{C \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D + 2\varepsilon \sqrt{CE} \coth\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{117}$$

$$v(x, t) = \chi \left[ \frac{C \operatorname{csch}^2\left(\frac{n}{2} \sqrt{C}(x - ct)\right)}{D + 2\varepsilon \sqrt{CE} \coth\left(\frac{n}{2} \sqrt{C}(x - ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{118}$$

provided  $\varepsilon = \pm 1$  and  $E > 0$ .

**(VIII)** If  $D^2 - 4CE > 0$  and  $C < 0$ , then one gets the periodic solutions:

$$v(x, t) = \left[ \frac{-2C \sec^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)}{2\sqrt{D^2-4CE} - \left[\sqrt{D^2-4CE}-D\right] \sec^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{119}$$

$$v(x, t) = \chi \left[ \frac{-2C \sec^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)}{2\sqrt{D^2-4CE} - \left[\sqrt{D^2-4CE}-D\right] \sec^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{120}$$

or

$$u(x, t) = \left[ \frac{2C \csc^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)}{2\sqrt{D^2-4CE} - \left[\sqrt{D^2-4CE}+D\right] \csc^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{121}$$

$$v(x, t) = \chi \left[ \frac{2C \csc^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)}{2\sqrt{D^2-4CE} - \left[\sqrt{D^2-4CE}+D\right] \csc^2\left(\frac{\sqrt{-C}}{2}n(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{122}$$

or

$$u(x, t) = \left[ \frac{2C \sec(n\sqrt{-C}(x-ct))}{\varepsilon\sqrt{D^2-4CE} - D \sec(n\sqrt{-C}(x-ct))} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{123}$$

$$v(x, t) = \chi \left[ \frac{2C \sec(n\sqrt{-C}(x-ct))}{\varepsilon\sqrt{D^2-4CE} - D \sec(n\sqrt{-C}(x-ct))} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{124}$$

or

$$u(x, t) = \left[ \frac{2C \csc(n\sqrt{-C}(x-ct))}{\varepsilon\sqrt{D^2-4CE} - D \csc(n\sqrt{-C}(x-ct))} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{125}$$

$$v(x, t) = \chi \left[ \frac{2C \csc(n\sqrt{-C}(x-ct))}{\varepsilon\sqrt{D^2-4CE} - D \csc(n\sqrt{-C}(x-ct))} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{126}$$

provided  $\varepsilon = \pm 1$ .

**(IX)** If  $C < 0$ , then one gets the periodic solutions:

$$u(x, t) = \left[ -\frac{C \sec^2\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)}{D + 2\varepsilon\sqrt{-CE} \tan\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{127}$$

$$v(x, t) = \chi \left[ -\frac{C \sec^2\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)}{D + 2\varepsilon\sqrt{-CE} \tan\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{128}$$

or

$$u(x, t) = \left[ -\frac{C \csc^2\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)}{D + 2\varepsilon\sqrt{-CE} \cot\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{129}$$

$$v(x, t) = \chi \left[ -\frac{C \csc^2\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)}{D + 2\varepsilon\sqrt{-CE} \cot\left(\frac{n}{2}\sqrt{-C}(x-ct)\right)} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{130}$$

provided  $\varepsilon = \pm 1$  and  $E > 0$ .

(X) If  $C > 0$ , then one gets the solutions:

$$u(x, t) = \left[ \frac{4Cn^2 e^{\varepsilon n \sqrt{C}(x-ct)}}{(e^{\varepsilon n \sqrt{C}(x-ct)} - Dn^2)^2 - 4CEn^4} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{131}$$

$$v(x, t) = \chi \left[ \frac{4Cn^2 e^{\varepsilon n \sqrt{C}(x-ct)}}{(e^{\varepsilon n \sqrt{C}(x-ct)} - Dn^2)^2 - 4CEn^4} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{132}$$

provided  $\varepsilon = \pm 1$ .

(XI) If  $C > 0$  and  $D = 0$ , then one gets the solutions:

$$u(x, t) = \left[ \frac{4Cn^2 e^{\varepsilon n \sqrt{C}(x-ct)}}{-1 + 4CEn^4 e^{2\varepsilon n \sqrt{C}(x-ct)}} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{133}$$

$$v(x, t) = \chi \left[ \frac{4Cn^2 e^{\varepsilon n \sqrt{C}(x-ct)}}{-1 + 4CEn^4 e^{2\varepsilon n \sqrt{C}(x-ct)}} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{134}$$

provided  $\varepsilon = \pm 1$ .

(XII) If  $C = 0$  and  $D = 0$ , then one gets the solutions:

$$u(x, t) = \left[ \frac{\varepsilon}{n\sqrt{E}(x-ct)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{135}$$

$$v(x, t) = \chi \left[ \frac{\varepsilon}{n\sqrt{E}(x-ct)} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{136}$$

provided  $E > 0$  and  $\varepsilon = \pm 1$ .

Case-7: If we substitute  $A = 0, B = \frac{8C^2}{27D}, E = \frac{D^2}{4C}$ , in the above conditions (43)–(51), then one gets the results:

$$\begin{aligned} H_1 &= -\left[ \frac{4}{27}(5n^2 + 4) + 1 \right] C^2 - H_0 C, \\ \Delta_1 &= -\frac{13}{18}(2+n)D \left[ C \left( n^2 + \frac{58}{39}n + \frac{58}{39} \right) + \frac{9}{13}H_0 \right], \\ \Delta_2 &= -(1+n) \frac{D^2}{4C} [C(7n^2 + 12n + 6) + H_0], \\ \Delta_3 &= -(3n+2)(2n+1)(1+n) \frac{D^3}{4C}, \quad \Delta_4 = -(3n+1)(2n+1)(1+n) \frac{D^4}{16C^2}, \\ \Gamma_1 &= \frac{4(n-2)C^2}{27D} [(n^2 - 2n + 2)C + H_0], \quad \Gamma_2 = \frac{16(n-1)(3n^2 - 8n + 4)C^4}{729D^2}, \quad \Gamma_3 = 0, \quad \Gamma_4 = 0. \end{aligned} \tag{137}$$

Under the constraint conditions (137), we have the solutions:

(I) If  $C < 0$ , then one gets the dark soliton solutions:

$$u(x, t) = \left[ -\frac{8C \tanh^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}}(x-ct) \right)}{3D \left[ 3 + \tanh^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}}(x-ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{138}$$

$$v(x, t) = \chi \left[ -\frac{8C \tanh^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}}(x-ct) \right)}{3D \left[ 3 + \tanh^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}}(x-ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{139}$$

and the singular soliton solutions:

$$u(x, t) = \left[ -\frac{8C \operatorname{coth}^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}} (x - ct) \right)}{3D \left[ 3 + \operatorname{coth}^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}} (x - ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{140}$$

$$v(x, t) = \chi \left[ -\frac{8C \operatorname{coth}^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}} (x - ct) \right)}{3D \left[ 3 + \operatorname{coth}^2 \left( \frac{n}{2} \sqrt{-\frac{C}{3}} (x - ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)}. \tag{141}$$

(II) If  $C > 0$ , then one gets the periodic solutions:

$$u(x, t) = \left[ \frac{8C \tan^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right)}{3D \left[ 3 - \tan^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{142}$$

$$v(x, t) = \chi \left[ \frac{8C \tan^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right)}{3D \left[ 3 - \tan^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{143}$$

or

$$u(x, t) = \left[ \frac{8C \cot^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right)}{3D \left[ 3 - \cot^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{144}$$

$$v(x, t) = \chi \left[ \frac{8C \cot^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right)}{3D \left[ 3 - \cot^2 \left( \frac{n}{2} \sqrt{\frac{C}{3}} (x - ct) \right) \right]} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)}. \tag{145}$$

Case-8: If we substitute  $A = 0$  in the above conditions (43)–(51), then one gets the results

$$\begin{aligned} H_1 &= -\frac{1}{2}(5n^2 + 4)DB - C(C + H_0), \\ \Delta_1 &= -(2 + n) \left\{ \left[ \left( 3BE + \frac{1}{2}DC \right) n^2 + (n + 1)(BE + DC) \right] + \frac{1}{2}DH_0 \right\}, \\ \Delta_2 &= -(1 + n) \left\{ \left[ n^2 \left( \frac{3}{4}D^2 + 4EC \right) + (2n + 1)(D^2 + 2EC) \right] + H_0E \right\}, \\ \Delta_3 &= -(3n + 2)(2n + 1)(1 + n)ED, \quad \Delta_4 = -(3n + 1)(2n + 1)(1 + n)E^2, \\ \Gamma_1 &= \frac{1}{2}(n - 2)B[(n^2 - 2n + 2)C + H_0], \\ \Gamma_2 &= \frac{1}{4}(n - 1)(3n^2 - 8n + 4)B^2, \quad \Gamma_3 = 0, \quad \Gamma_4 = 0, \end{aligned} \tag{146}$$

Under the constraint conditions (146), we have the solutions:

(I) If  $B = \frac{D^3(m^2 - 1)}{32m^2E^2}$  and  $C = \frac{D^2(5m^2 - 1)}{16m^2E}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \operatorname{sn} \left( \frac{nD}{4m} \sqrt{\frac{1}{E}} (x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{147}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \varepsilon \operatorname{sn} \left( \frac{nD}{4m} \sqrt{\frac{1}{E}} (x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{148}$$

or

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{msn} \left( \frac{nD}{4m} \sqrt{\frac{1}{E}} (x - ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{149}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{msn} \left( \frac{nD}{4m} \sqrt{\frac{1}{E}} (x - ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{150}$$

provided  $\varepsilon = \pm 1$  and  $E > 0$ .

**(II)** If  $B = \frac{D^3(1-m^2)}{32E^2}$  and  $C = \frac{D^2(5-m^2)}{16E}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \varepsilon \operatorname{msn} \left( \frac{nD}{4} \sqrt{\frac{1}{E}} (x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{151}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \varepsilon \operatorname{msn} \left( \frac{nD}{4} \sqrt{\frac{1}{E}} (x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{152}$$

or

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{sn} \left( \frac{nD}{4} \sqrt{\frac{1}{E}} (x - ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{153}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{sn} \left( \frac{nD}{4} \sqrt{\frac{1}{E}} (x - ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{154}$$

provided  $\varepsilon = \pm 1$  and  $E > 0$ .

**(III)** If  $B = \frac{D^3}{32m^2E^2}$  and  $C = \frac{D^2(4m^2+1)}{16m^2E}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \varepsilon \operatorname{cn} \left( \frac{nD}{4m} \sqrt{-\frac{1}{E}} (x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{155}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \varepsilon \operatorname{cn} \left( \frac{nD}{4m} \sqrt{-\frac{1}{E}} (x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{156}$$

or

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon \sqrt{1-m^2} \operatorname{sn} \left( \frac{nD}{4m} \sqrt{-\frac{1}{E}} (x - ct) \right)}{\operatorname{dn} \left( \frac{nD}{4m} \sqrt{-\frac{1}{E}} (x - ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{157}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon \sqrt{1-m^2} \operatorname{sn} \left( \frac{nD}{4m} \sqrt{-\frac{1}{E}}(x-ct) \right)}{\operatorname{dn} \left( \frac{nD}{4m} \sqrt{-\frac{1}{E}}(x-ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{158}$$

provided  $\varepsilon = \pm 1$  and  $E < 0$ .

(IV) If  $B = \frac{m^2 D^3}{32E^2(m^2-1)}$  and  $C = \frac{D^2(5m^2-4)}{16E(m^2-1)}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\sqrt{1-m^2}} \operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{(1-m^2)E}}(x-ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{159}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\sqrt{1-m^2}} \operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{(1-m^2)E}}(x-ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{160}$$

or

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{(1-m^2)E}}(x-ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{161}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{(1-m^2)E}}(x-ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{162}$$

provided  $\varepsilon = \pm 1$  and  $E < 0$ .

(VI) If  $B = \frac{D^3}{32E^2(1-m^2)}$  and  $C = \frac{D^2(4m^2-5)}{16E(m^2-1)}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{cn} \left( \frac{nD}{4} \sqrt{\frac{1}{(1-m^2)E}}(x-ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{163}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon}{\operatorname{cn} \left( \frac{nD}{4} \sqrt{\frac{1}{(1-m^2)E}}(x-ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{164}$$

or

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon \operatorname{dn} \left( \frac{nD}{4} \sqrt{\frac{1}{(1-m^2)E}}(x-ct) \right)}{\sqrt{1-m^2} \operatorname{sn} \left( \frac{nD}{4} \sqrt{\frac{1}{(1-m^2)E}}(x-ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{165}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon \operatorname{dn} \left( \frac{nD}{4} \sqrt{\frac{1}{(1-m^2)E}}(x-ct) \right)}{\sqrt{1-m^2} \operatorname{sn} \left( \frac{nD}{4} \sqrt{\frac{1}{(1-m^2)E}}(x-ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{166}$$

provided  $\varepsilon = \pm 1$  and  $E > 0$ .

(VII) If  $B = \frac{m^2 D^3}{32E^2}$  and  $C = \frac{D^2(m^2+4)}{16E}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:



$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{E}}(x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{167}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{E}}(x - ct) \right) \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{168}$$

or

$$u(x, t) = \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon \sqrt{1 - m^2}}{\operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{E}}(x - ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{169}$$

$$v(x, t) = \chi \left[ -\frac{D}{4E} \left\{ 1 + \frac{\varepsilon \sqrt{1 - m^2}}{\operatorname{dn} \left( \frac{nD}{4} \sqrt{-\frac{1}{E}}(x - ct) \right)} \right\} \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{170}$$

provided  $\varepsilon = \pm 1$  and  $E < 0$ .

**Case-9:** If we substitute  $A = E = 0$  in the above conditions (43)–(51), then one gets the results:

$$\begin{aligned} H_1 &= - \left[ \left( \frac{5}{2}n^2 + 2 \right) DB + C^2 + H_0 C \right], \\ \Delta_1 &= -(2 + n)D \{ (n^2 + 2n + 2)C + H_0 \}, \quad \Delta_2 = -(1 + n) \left( \frac{3}{4}n^2 + 2n + 1 \right) D^2, \\ \Delta_3 &= 0, \quad \Delta_4 = 0, \quad \Gamma_1 = \frac{(n - 2)}{2} B [ (n^2 - 2n + 2)C + H_0 ], \\ \Gamma_2 &= (n - 1) \left( \frac{3}{4}n^2 - 2n + 1 \right) B^2, \quad \Gamma_3 = 0, \quad \Gamma_4 = 0. \end{aligned} \tag{171}$$

Under the constraint conditions (171), we have the solutions:

(I) If  $B = \frac{C^2 m^2 (m^2 - 1)}{D(2m^2 - 1)^2}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{m^2 C}{D(2m^2 - 1)} \operatorname{cn}^2 \left( \frac{n}{2} \sqrt{\frac{C}{2m^2 - 1}}(x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{172}$$

$$v(x, t) = \left[ -\frac{m^2 C}{D(2m^2 - 1)} \operatorname{cn}^2 \left( \frac{n}{2} \sqrt{\frac{C}{2m^2 - 1}}(x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{173}$$

provided  $D < 0$  and  $C(2m^2 - 1) > 0$ .

(II) If  $B = \frac{C^2 m^2}{D(m^2 + 1)^2}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{m^2 C}{D(m^2 + 1)} \operatorname{sn}^2 \left( \frac{n}{2} \sqrt{-\frac{C}{m^2 + 1}}(x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{174}$$

$$v(x, t) = \chi \left[ -\frac{m^2 C}{D(m^2 + 1)} \operatorname{sn}^2 \left( \frac{n}{2} \sqrt{-\frac{C}{m^2 + 1}}(x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{175}$$

or

$$u(x, t) = \left[ -\frac{m^2 C}{D(m^2 + 1)} \operatorname{cd}^2 \left( \frac{n}{2} \sqrt{\frac{C}{m^2 + 1}}(x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{176}$$

$$v(x, t) = \chi \left[ -\frac{m^2 C}{D(m^2 + 1)} \operatorname{cd}^2 \left( \frac{n}{2} \sqrt{\frac{C}{m^2 + 1}}(x - ct) \right) \right]^{\frac{1}{n}} e^{i(-kx + \omega t + \theta_0)} \tag{177}$$

provided  $D > 0$  and  $C < 0$ .

(III) If  $B = \frac{C^2(1-m^2)}{D(2-m^2)^2}$  where  $0 < m < 1$ , then one gets the Jacobi elliptic function solutions:

$$u(x, t) = \left[ -\frac{C}{D(2-m^2)} \operatorname{dn}^2 \left( \frac{n}{2} \sqrt{\frac{C}{2-m^2}} (x-ct) \right) \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{178}$$

$$v(x, t) = \chi \left[ -\frac{C}{D(2-m^2)} \operatorname{dn}^2 \left( \frac{n}{2} \sqrt{\frac{C}{2-m^2}} (x-ct) \right) \right]^{\frac{1}{n}} e^{i(-kx+\omega t+\theta_0)} \tag{179}$$

provided  $D < 0$  and  $C > 0$ . Finally, if  $m \rightarrow 1^-$  or  $m \rightarrow 0^+$  in the above Jacobi elliptic function solutions, we can get the hyperbolic and periodic solutions, respectively.

#### 4. Conservation laws

In the system (2)-(3) above, we let  $u = U + iW$  and  $v = V + iZ$  so that the system splits into a system of four pdes whose conserved flows ( $T^t, T^x$ ) are constructed using the multiplier approach. It turns out that if  $Q_1 = Q_2$ , we have a single multiplier  $Q = (-u, -v, w, z)$  giving rise to the conserved density

$$T_1^t = \frac{1}{2} (U^2 + W^2 + V^2 + Z^2) \tag{180}$$

so that a corresponding conserved density of the complex system (2)-(3) is

$$\Phi_1^t = |u|^2 + |v|^2.$$

Unless,  $\tau_i = 0, \zeta_i = 0, \eta_i = 0, \xi_i = 0, \alpha_i = 0, \beta_i = 0, \gamma_i = 0, \delta_i = 0$  and  $\nu_i = -\lambda_i$ , for  $i = 1, 2$ , No other conservation laws exist. Under these conditions, we have momentum and energy conserved.

The following conserved density,  $T^t$ , for linear momentum is

$$T_2^t = - \left[ -\frac{1}{2} V_x Z - \frac{1}{2} U_x W + \frac{1}{2} Z_x U + \frac{1}{2} W_x V \right] \tag{181}$$

and the momentum density is

$$\Phi_2^t = \mathcal{I}(u^* u_x) + \mathcal{I}(v^* v_x). \tag{182}$$

Some of the terms in the conserved density corresponding to Hamiltonian are

$$\begin{aligned} T_3^t = & 169 (V^2 + Z^2)^n V^2 f_2 n^4 + 36 (V^2 + Z^2)^n Z^2 f_2 n^7 - 16 (V^2 + Z^2)^n Z^2 f_2 n - 36 (V^2 + Z^2)^n V^2 f_2 n^6 \\ & - 36 (V^2 + Z^2)^n Z^2 f_2 n^6 - 169 (V^2 + Z^2)^n V^2 f_2 n^5 - 169 (V^2 + Z^2)^n Z^2 f_2 n^5 - 98 (V^2 + Z^2)^{2n} V^2 h_2 n^5 \\ & - 9 (V^2 + Z^2)^{2n} Z^2 h_2 n^6 - 9 (V^2 + Z^2)^{2n} V^2 h_2 n^6 + 18 (V^2 + Z^2)^{2n} Z^2 h_2 n^7 + 18 (V^2 + Z^2)^{2n} V^2 h_2 n^7 \\ & - 32 (V^2 + Z^2)^{2n} Z^2 h_2 n - 32 (V^2 + Z^2)^{2n} V^2 h_2 n - 56 (V^2 + Z^2)^{2n} Z^2 h_2 n^2 + 49 (V^2 + Z^2)^{2n} Z^2 h_2 n^4 \\ & \dots \end{aligned} \tag{183}$$

It is clear that the density is not only cumbersome but when written in terms of  $u$  and  $v$  would be divergent.

The bright solitons along the two components of the magneto-optic waveguides take the form:

$$u(x, t) = \frac{A_1}{\{D + \cosh[B(x-ct)]\}^{\frac{1}{n}}} e^{i(-kx+\omega t+\theta_0)} \tag{184}$$

and

$$v(x, t) = \frac{A_2}{\{D + \cosh[B(x-ct)]\}^{\frac{1}{n}}} e^{i(-kx+\omega t+\theta_0)}. \tag{185}$$

Thus, for these bright solitons, the two conserved quantities would be

$$P = \int_{-\infty}^{\infty} (|u|^2 + |v|^2) dx = \frac{2}{2^n B} (A_1^2 + A_2^2) F \left( \frac{2}{n}, \frac{2}{n}; \frac{2}{n} + \frac{1}{2}; \frac{1-D}{2} \right) \frac{\Gamma \left( \frac{2}{n} \right) \left( \frac{1}{2} \right)}{\Gamma \left( \frac{2}{n} + \frac{1}{2} \right)} \tag{186}$$

and

$$M = i \int_{-\infty}^{\infty} \{ (uu_x^* - u^*u_x) + (vv_x^* - v^*v_x) \} dx = \frac{2\kappa}{2^{\frac{1}{2}}B} (A_1^2 + A_2^2) F\left(\frac{2}{n}; \frac{2}{n}; \frac{2}{n} + \frac{1}{2}; \frac{1-D}{2}\right) \frac{\Gamma\left(\frac{2}{n}\right)\Gamma\left(\frac{1}{2}\right)}{\Gamma\left(\frac{2}{n} + \frac{1}{2}\right)} \quad (187)$$

which represent total power and linear momentum of the solitons, respectively. Here, Gauss' hypergeometric function is written as:

$$F(\alpha, \beta; \gamma; z) = \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n} \frac{z^n}{n!} \quad (188)$$

where the Pochhammer symbol is:

$$(p)_n = \begin{cases} 1 & n = 0, \\ p(p+1)\cdots(p+n-1) & n > 0. \end{cases} \quad (189)$$

The condition that guarantees convergence of the series is

$$|z| < 1 \quad (190)$$

which for (186) and (187) means

$$-1 < D < 3. \quad (191)$$

Finally, Rabbe's test of convergence reveals another criterion for convergence of the series, namely

$$\gamma < \alpha + \beta \quad (192)$$

which transforms to

$$0 < n < 4. \quad (193)$$

This gives the zone of existence of the solitons which simply means the generalized KE will yield solitons in magneto-optic waveguides provided (193) stays valid. Benjamin-Fier stability analysis would also yield the same condition for the existence of solitons.

## 5. Conclusions

This paper revealed soliton solutions in terms of generalized KE that are recovered through an intermediate layer of solutions, namely Jacobi's elliptic functions. The limiting process eventually revealed the soliton solutions that are classified as bright, dark, singular types. Another class of solutions were also recovered, and those are in terms of Weierstrass' elliptic functions. Finally, the conservation laws are formulated. The power and linear momentum are the only two conservation laws that fall out of the model. A natural consequence of conservation laws is the stability zone for the solitons that naturally emerged from the convergence criteria of Gauss' hypergeometric functions that yielded bounds for the power law parameter  $n$ . This therefore avoided the computation of the stability zone for the solitons by the usage of Benjamin-Fier stability analysis. Pretty slick!

Later, further research activities will be carried out to secure furthermore novel and interesting results to the model utilizing a variety of integration schemes that have been successfully performed in this field of research [26–45]. Studies in this context are in progress and will be reported with time.

## Declaration of Competing Interest

The authors also declare that there is no conflict of interest.

## Acknowledgments

The research work of the ninth author (MRB) was supported by the grant NPRP 11S-1126-170033 from QNRF and he is thankful for it.

## References

- [1] M. Asma, A. Biswas, A.H. Kara, E.M.E. Zayed, P. Guggilla, S. Khan, M. Ekici, A.K. Alzahrani, M.R. Belic, A pen-picture of solitons and conservation laws in magneto-optic waveguides having quadratic-cubic law of nonlinear refractive index, *Optik* 223 (2020) 165330.
- [2] A. Biswas, A.H. Arnous, M. Ekici, A. Sonmezoglu, A.R. Seadawy, Q. Zhou, M.F. Mahmood, S.P. Moshokoa, M. Belic, Optical soliton perturbation in magneto-optic waveguides, *J. Nonlinear Opt. Phys. Mater.* 27 (2018) 1850005.
- [3] A. Biswas, Solitons in magneto-optic waveguides, *Appl. Math. Comput.* 153 (2004) 387–393.

- [4] J.V. Guzman, A.A. Alshaery, E.M. Hilal, A.H. Bhrawy, M.F. Mahmood, L. Moraru, A. Biswas, Optical soliton perturbation in magneto-optic waveguides with spatiotemporal dispersion, *J. Optoelectron. Adv. Mater.* 16 (9–10) (2014) 1063–1070.
- [5] H. Dötsch, N. Bahlmann, O. Zhuromskyy, M. Hammer, L. Wilkens, R. Gerhardt, P. Hertel, A.F. Popkov, Applications of magneto-optical waveguides in integrated optics: review, *J. Opt. Soc. Am. B* 22 (1) (2005) 240–253.
- [6] T. Haider, A review of magneto-optic effects and its application, *Int. J. Electromagn. Appl.* 7 (1) (2017) 17–24.
- [7] K. Hasegawa, Y. Miyazaki, Magneto-optic devices using interaction between magnetostatic surface wave and optical guided wave, *Jpn. J. Appl. Phys.* 31 (1992) 230.
- [8] A.H. Kara, A. Biswas, M. Belic, Conservation laws for optical solitons in birefringent fibers and magneto-optic waveguides, *Optik* 127 (2016) 11662–11673.
- [9] N.A. Kudryashov, A generalized model for description of propagation pulses in optical fiber, *Optik* 189 (2019) 42–52.
- [10] N.A. Kudryashov, Mathematical model of propagation pulse in optical fiber with power nonlinearities, *Optik* 212 (2020) 164750.
- [11] N.A. Kudryashov, E.V. Antonova, Solitary waves of equation for propagation pulse with power nonlinearities, *Optik* 217 (2020) 164881.
- [12] N.A. Kudryashov, Periodic and solitary waves in optical fiber Bragg gratings with dispersive reflectivity, *Chin. J. Phys.* 66 (2020) 401–405.
- [13] Y. Qiu, B.A. Malomed, D. Mihalache, X. Zhu, J. Peng, Y. He, Generation of stable multi-vortex clusters in a dissipative medium with anti-cubic nonlinearity, *Phys. Lett. A* 383 (228) (2019) 2579–2583.
- [14] Y. Shoji, T. Mizumoto, Waveguide magneto-optical devices for photonics integrated circuits, *Opt. Mater. Express* 8 (8) (2018) 2387–2394.
- [15] E.M.E. Zayed, R.M.A. Shohib, M.M. El-Horbaty, A. Biswas, M. Asma, M. Ekici, A.K. Alzahrani, M.R. Belic, Solitons in magneto-optic waveguides with quadratic-cubic nonlinearity, *Phys. Lett. A* 384 (25) (2020) 126456.
- [16] E.M.E. Zayed, R.M.A. Shohib, M.E.M. Alngar, A. Biswas, M. Asma, M. Ekici, S.P. Moshokoa, A.K. Alzahrani, M.R. Belic, Solitons in magneto-optic waveguides with dual-power law nonlinearity, *Phys. Lett. A* 384 (27) (2020) 126697.
- [17] E.M.E. Zayed, M.E.M. Alngar, R.M.A. Shohib, A. Biswas, M. Ekici, A.K. Alzahrani, M.R. Belic, Solitons in magneto-optic waveguides with anti-cubic nonlinearity, *Optik* 222 (2020) 165313.
- [18] E.M.E. Zayed, M.E.M. Alngar, M.M. El-Horbaty, A. Biswas, M. Asma, M. Ekici, A.K. Alzahrani, M.R. Belic, Solitons in magneto-optic waveguides with generalized anti-cubic nonlinearity, *Optik* 223 (2020) 165456.
- [19] E.M.E. Zayed, M.E.M. Alngar, A. Biswas, M. Asma, M. Ekici, A.K. Alzahrani, M.R. Belic, Solitons in magneto-optic waveguides with Kudryashov's law of refractive index, *Chaos Solitons Fractals* 140 (2020) 110129.
- [20] E.M.E. Zayed, M.E.M. Alngar, Application of newly proposed sub-ODE method to locate chirped optical solitons to Triki–Biswas equation, *Optik* 207 (2020) 164360.
- [21] Z. Li-Hua, H. Jin-Yu, Sub-ODE's new solutions and their applications to two nonlinear partial differential equations with higher-order nonlinear terms, *Commun. Theor. Phys.* 52 (5) (2009) 773–778.
- [22] Z.L. Li, Periodic wave solutions of a generalized KdV-mKdV equation with higher-order nonlinear terms, *Z. Naturforsch.* 56a (2010) 649–657.
- [23] S. El-Ganaini, H. Kumar, A variety of new traveling and localized solitary wave solutions of a nonlinear model describing the nonlinear low-pass electrical transmission lines, *Chaos Solitons Fractals* 140 (2020) 110218.
- [24] E.M.E. Zayed, M.E.M. Alngar, A. Biswas, H. Triki, Y.Y.r. m, A.S. Alshomrani, Chirped and chirp-free optical solitons in fiber Bragg gratings with dispersive reflectivity having quadratic-cubic nonlinearity by sub-ODE approach, *Optik* 203 (2020) 163993.
- [25] E.M.E. Zayed, M.E.M. Alngar, A. Biswas, M. Ekici, H. Triki, A.K. Alzahrani, M.R. Belic, Chirped and chirp-free optical solitons in fiber Bragg gratings having dispersive reflectivity with polynomial form of nonlinearity using sub-ODE approach, *Optik* 204 (2020) 164096.
- [26] N.A. Kudryashov, Polynomials in logistic function and solitary waves of nonlinear differential equations, *Appl. Math. Comput.* 219 (17) (2013) 9245–9253.
- [27] N.A. Kudryashov, Logistic function as solution of many nonlinear differential equations, *Appl. Math. Model.* 39 (18) (2015) 5733–5742.
- [28] N.A. Kudryashov, Method for finding highly dispersive optical solitons of nonlinear differential equations, *Optik* 206 (2020) 163550.
- [29] N.A. Kudryashov, Highly dispersive optical solitons of equation with various polynomial nonlinearity law, *Chaos Solitons Fractals* 140 (2020) 110202.
- [30] N.A. Kudryashov, Highly dispersive solitary wave solutions of perturbed nonlinear Schrödinger equations, *Appl. Math. Comput.* 371 (2020) 124972.
- [31] N.A. Kudryashov, Solitary wave solutions of hierarchy with non-local nonlinearity, *Appl. Math. Lett.* 103 (2020) 106155.
- [32] N.A. Kudryashov, The generalized duffing oscillator, *Commun. Nonlinear Sci. Numer. Simul.* 93 (2021) 105526.
- [33] J. Dan, S. Sain, A. Ghose-Choudhury, S. Garai, Application of the Kudryashov function for finding solitary wave solutions of NLS type differential equations, *Optik* 224 (2020) 165519.
- [34] Y. Han, B. Tian, Y.Q. Yuan, C.R. Zhang, S.S. Chen, Bilinear forms and bright-dark solitons for a coupled nonlinear Schrödinger system with variable coefficients in an inhomogeneous optical fiber, *Chin. J. Phys.* 62 (2019) 202–212.
- [35] A.M. Wazwaz, Multiple complex soliton solutions for the integrable KdV, fifth-order Lax, modified KdV, Burgers, and Sharma–Tasso–Olver equations, *Chin. J. Phys.* 59 (2019) 372–378.
- [36] A.M. Wazwaz, The integrable Vakhnenko–Parkes (VP) and the modified Vakhnenko–Parkes (MVP) equations: multiple real and complex soliton solutions, *Chin. J. Phys.* 57 (2019) 375–381.
- [37] A.H. Arnous, M. Ekici, A. Biswas, A.S. Alshomrani, M.R. Belic, Optical solitons having anti-cubic nonlinearity with two integration architectures, *Chin. J. Phys.* 60 (2019) 659–664.
- [38] A. Biswas, C.M. Khalique, Stationary solutions for the nonlinear dispersive Schrödinger equation with generalized evolution, *Chin. J. Phys.* 51 (1) (2013) 103–110.
- [39] A. Das, A. Biswas, M. Ekici, Q. Zhou, M.R. Belic, Optical solitons with complex Ginzburg–Landau equation for two nonlinear forms using  $F$ -expansion, *Chin. J. Phys.* 61 (2019) 255–261.
- [40] E.V. Krishnan, A. Biswas, Q. Zhou, M. Ekici, A.S. Alshomrani, M. Belic, Optical soliton perturbation with quadratic-cubic nonlinearity by mapping methods, *Chin. J. Phys.* 60 (2019) 632–637.
- [41] X. Liu, Z. Luan, Q. Zhou, W. Liu, A. Biswas, Dark two-soliton solutions for nonlinear Schrödinger equations in inhomogeneous optical fibers, *Chin. J. Phys.* 61 (2019) 310–315.
- [42] J.-Y. Song, Y.K. Yang, Soliton solutions and conservation laws for a generalized Ablowitz–Ladik system, *Chin. J. Phys.* 60 (2019) 271–278.
- [43] X.-Y. Xie, G.Q. Meng, Dark soliton excitations and collisions for the  $(2+1)$ -dimensional variable-coefficient Davey–Stewartson-like equations in the plasmas or Bose–Einstein condensates, *Chin. J. Phys.* 59 (2019) 160–165.
- [44] Y.Y.l.r. m, M. Mirzazadeh, Optical pulses with Kundu–Mukherjee–Naskar model in fiber communication systems, *Chin. J. Phys.* 64 (2020) 183–193.
- [45] Y.Y. Yıldırım, A. Biswas, M. Ekici, O. Gonzalez-Gaxiola, S. Khan, H. Triki, L. Moraru, A.K. Alzahrani, M.R. Belic, Optical solitons with Kudryashov's model by a range of integration norms, *Chin. J. Phys.* 66 (2020) 660–672.