

## Cubic–quartic polarized optical solitons and conservation laws for perturbed Fokas–Lenells model

Elsayed M. E. Zayed<sup>\*</sup>, Mohamed E. M. Alngar<sup>\*</sup>, Mahmoud M. El-Horbaty<sup>\*</sup>,  
Anjan Biswas<sup>†,‡,§,¶</sup>, Abdul H. Kara<sup>||</sup>, Yakup Yildirim<sup>\*\*,\*\*,‡‡</sup>, Salam Khan<sup>¶</sup>,  
Abdullah Khamis Alzahrani<sup>‡</sup> and Milivoj R. Belic<sup>††</sup>

<sup>\*</sup>Mathematics Department, Faculty of Science,  
Zagazig University, Zagazig 44519, Egypt

<sup>†</sup>Department of Applied Mathematics,  
National Research Nuclear University,  
31 Kashirskoe Hwy, Moscow 115409, Russian Federation

<sup>‡</sup>Mathematical Modeling and Applied Computation (MMAC) Research Group,  
Department of Mathematics, King Abdulaziz University,  
Jeddah 21589, Saudi Arabia

<sup>§</sup>Department of Mathematics and Applied Mathematics,  
Sefako Makgatho Health Sciences University,  
Medunsa 0204, Pretoria, South Africa

<sup>¶</sup>Department of Physics, Chemistry and Mathematics,  
Alabama A&M University, Normal, AL 35762-4900, USA

<sup>||</sup>School of Mathematics, University of the Witwatersrand,  
Private Bag 3, Wits-2050, Johannesburg, South Africa

<sup>\*\*</sup>Department of Mathematics, Faculty of Arts and Sciences,  
Near East University, 99138 Nicosia, Cyprus

<sup>††</sup>Institute of Physics Belgrade, Pregrevice 118,  
11080 Zemun, Serbia

<sup>\*\*</sup>yakup.yildirim@neu.edu.tr

Received 6 February 2021

Revised 30 March 2021

Accepted 16 July 2021

Published 8 September 2021

This paper studies polarized cubic–quartic solitons that are modeled by Fokas–Lenells equation in birefringent fibers. Two integration schemes recovered a spectrum of soliton solutions to the model. Subsequently, the bright solitons compute the corresponding conserved quantities from the respective densities that are recovered by the multiplier approach.

*Keywords:* Solitons; polarization; Fokas–Lenells.

OCIS Codes: 060.2310, 060.4510, 060.5530, 190.3270, 190.4370.

<sup>‡‡</sup>Corresponding author.

## 1. Introduction

The concept of cubic–quartic (CQ) optical solitons first appeared a couple of years ago as a generalization to pure-quartic solitons and pure-cubic solitons that were introduced to sustain the delicate balance between chromatic dispersion (CD) and nonlinearity. The issue with pure-quartic solitons is that an analytical soliton solution cannot be established unless one is looking for a stationary soliton solution. On the other hand, pure-cubic soliton model is not integrable unless Hamiltonian perturbation terms are included in it. This led to the novel concept of CQ solitons that has achieved success in the fields of quantum optics and telecommunication industry. In CQ solitons, CD is replaced by a combination of third-order dispersion (3OD) and fourth-order dispersion (4OD) that can together replenish the low count of CD. A plethora of results are reported during the past decade or so in these arenas.<sup>1–27</sup>

One of the familiar models that addresses the dynamics of soliton propagation through optical fibers and PCF, apart from the familiar nonlinear Schrödinger’s equation, is Fokas–Lenells equation (FLE). This model is well studied in the context of polarization-preserving fibers with CD and/or CQ replacing CD. Today’s work turns the page to extend the study of CQ solitons with FLE for polarized pulses. Thus, the model equations for CQ solitons in birefringent fibers are first introduced. Subsequently, a couple of familiar integration algorithms would give way to a spectrum of CQ soliton solutions to FLE in birefringent fibers. Finally, the paper enumerates and exhibits conservation laws to the model that are recovered by the aid of multiplier approach. The corresponding conserved quantities are eventually computed and presented. These results are displayed after a succinct intro to the governing model.

### 1.1. Governing model

The CQ in birefringent fibers for the perturbed FLE reads as follows:

$$\begin{aligned}
 iq_t + ia_1q_{xxx} + b_1q_{xxxx} + (c_1|q|^2 + d_1|r|^2)(e_1q + if_1q_x) + qr^*(\gamma_1r + i\eta_1r_x) \\
 = i[\delta_1q_x + \lambda_1(|q|^2q)_x + \mu_1(|q|^2_xq)],
 \end{aligned}
 \tag{1}$$

$$\begin{aligned}
 ir_t + ia_2r_{xxx} + b_2r_{xxxx} + (c_2|r|^2 + d_2|q|^2)(e_2r + if_2r_x) + rq^*(\gamma_2q + i\eta_2q_x) \\
 = i[\delta_2r_x + \lambda_2(|r|^2r)_x + \mu_2(|r|^2_xr)],
 \end{aligned}
 \tag{2}$$

where  $a_j, b_j, c_j, d_j, e_j, f_j, \gamma_j, \eta_j, \delta_j, \lambda_j$  and  $\mu_j$  for  $(j = 1, 2)$  are all real-valued constants and  $i = \sqrt{-1}$ , while the independent variables  $x$  and  $t$  represent spatial and temporal variables, respectively. The dependent variables  $q = q(x, t)$  and  $r = r(x, t)$  are the complex valued describing the wave profiles for the two components in polarization-preserving fibers. The first terms in Eqs. (1) and (2) are linear temporal evolution of the pulses. Next, the coefficients of  $a_j$  are the 3ODs and the coefficients  $b_j$

stand for the 4ODs. Then,  $c_j$  are the self-phase modulation (SPM) terms while  $d_j$  stand for cross-phase modulation (XPM). Next,  $f_j$  represent nonlinear dispersions, while  $\eta_j$  are four wave-mixing (4WM) terms. The parameters  $e_j$  coupled with  $c_j$  give SPM and when coupled with  $d_j$  give XPM. Then  $\gamma_j$  give the XPM along with 4WM. Finally,  $\delta_j$  are the coefficients of inter-modal dispersion (IMD) and the parameters  $\lambda_j$  give self-steepening terms (SSTs) that avoid the formation of shock waves, while  $\mu_j$  are nonlinear dispersion coefficients.

This paper is organized as follows. In Sec. 2, the mathematical preliminaries are introduced. In Secs. 3 and 4, the generalized auxiliary equation method and the addendum to Kudryashov’s method are successfully applied. In Sec. 6, the conservation laws of the model are addressed with a few conclusive words that are given in Sec. 7.

## 2. Mathematical Preliminaries

In order to solve Eqs. (1) and (2), we assume the hypothesis:

$$\begin{aligned} q(x, t) &= U_1(\xi)\exp[i\psi(x, t)], \\ r(x, t) &= U_2(\xi)\exp[i\psi(x, t)], \end{aligned} \tag{3}$$

such that

$$\xi = x - vt, \quad \psi(x, t) = -\kappa x + \omega t + \theta_0, \tag{4}$$

where  $v, \kappa, \omega$  and  $\theta_0$  are all nonzero constants to be determined which represent soliton velocity, soliton frequency, wave number and phase constant, respectively. Next,  $\psi(x, t)$  is a real function which represents the phase component of the soliton, while  $U_j(\xi)$  are real functions which represent the shape of the pulse of the solitons. Substituting (3) along with (4) into Eqs. (1) and (2), separating the real and imaginary parts, we deduce that the real parts are

$$\begin{aligned} b_1 U_1'''' + (3a_1\kappa - 6b_1\kappa^2)U_1'' - [\omega + \delta_1\kappa + a_1\kappa^3 - b_1\kappa^4]U_1 \\ + (e_1 + f_1\kappa)(c_1U_1^2 + d_1U_2^2)U_1 + (\gamma_1 + \eta_1\kappa)U_1U_2^2 - \lambda_1\kappa U_1^3 = 0 \end{aligned} \tag{5}$$

and

$$\begin{aligned} b_2 U_2'''' + (3a_2\kappa - 6b_2\kappa^2)U_2'' - [\omega + \delta_2\kappa + a_2\kappa^3 - b_2\kappa^4]U_2 \\ + (e_2 + f_2\kappa)(c_2U_2^2 + d_2U_1^2)U_2 + (\gamma_2 + \eta_2\kappa)U_2U_1^2 - \lambda_2\kappa U_2^3 = 0, \end{aligned} \tag{6}$$

while the imaginary parts are

$$\begin{aligned} (a_1 - 4b_1\kappa)U_1''' - (\delta_1 + v + 3a_1\kappa^2 - 4b_1\kappa^3)U_1' - (3\lambda_1 + 2\mu_1)U_1^2U_1' \\ + f_1(c_1U_1^2 + d_1U_2^2)U_1' + \eta_1U_1U_2U_2' = 0 \end{aligned} \tag{7}$$

and

$$(a_2 - 4b_2\kappa)U_2''' - (\delta_2 + v + 3a_2\kappa^2 - 4b_2\kappa^3)U_2' - (3\lambda_2 + 2\mu_2)U_2^2U_2' + f_2(c_2U_2^2 + d_2U_1^2)U_2' + \eta_2U_2U_1U_1' = 0. \tag{8}$$

Set

$$U_2(\xi) = \chi U_1(\xi), \tag{9}$$

where  $\chi$  is a nonzero constant, such that  $\chi \neq 1$ . Consequently, the real parts change to

$$b_1U_1'''' + (3a_1\kappa - 6b_1\kappa^2)U_1'' - [\omega + \delta_1\kappa + a_1\kappa^3 - b_1\kappa^4]U_1 + [(e_1 + f_1\kappa)(c_1 + d_1\chi^2) + (\gamma_1 + \eta_1\kappa)\chi^2 - \lambda_1\kappa]U_1^3 = 0 \tag{10}$$

and

$$\chi b_2U_1'''' + (3a_2\kappa - 6b_2\kappa^2)\chi U_1'' - [\omega + \delta_2\kappa + a_2\kappa^3 - b_2\kappa^4]\chi U_1 + [(e_2 + f_2\kappa)(c_2\chi^2 + d_2)\chi + (\gamma_2 + \eta_2\kappa)\chi - \lambda_2\kappa\chi^3]U_1^3 = 0, \tag{11}$$

while the imaginary parts become

$$(a_1 - 4b_1\kappa)U_1''' - (\delta_1 + v + 3a_1\kappa^2 - 4b_1\kappa^3)U_1' + [f_1(c_1 + d_1\chi^2) - (3\lambda_1 + 2\mu_1) + \eta_1\chi^2]U_1^2U_1' = 0 \tag{12}$$

and

$$(a_2 - 4b_2\kappa)\chi U_1''' - (\delta_2 + v + 3a_2\kappa^2 - 4b_2\kappa^3)\chi U_1' + [f_2(c_2\chi^2 + d_2)\chi - (3\lambda_2 + 2\mu_2)\chi^3 + \eta_2\chi]U_1^2U_1' = 0. \tag{13}$$

The linearly independent principle is applied on (12) and (13) to get the frequency of soliton:

$$\kappa = \frac{a_j}{4b_j}, \quad a_j \neq 0, \quad b_j \neq 0 \quad \text{for } j = 1, 2, \tag{14}$$

the velocity of soliton is given by

$$v = 4b_j\kappa^3 - \delta_j - 3a_j\kappa^2 \quad \text{for } j = 1, 2, \tag{15}$$

and the constraint conditions

$$f_1(c_1 + d_1\chi^2) - (3\lambda_1 + 2\mu_1) + \eta_1\chi^2 = 0, \tag{16}$$

$$f_2(c_2\chi^2 + d_2) - (3\lambda_2 + 2\mu_2)\chi^2 + \eta_2 = 0.$$

From (14) and (15), one gets the natural constraint relations as

$$a_1b_2 = a_2b_1, \tag{17}$$

$$4b_1\kappa^3 - \delta_1 - 3a_1\kappa^2 = 4b_2\kappa^3 - \delta_2 - 3a_2\kappa^2.$$

Equations (10) and (11) have the same form under the constraint conditions:

$$\begin{aligned}
 b_1 &= \chi b_2, \\
 3a_1\kappa - 6b_1\kappa^2 &= (3a_2\kappa - 6b_2\kappa^2)\chi, \\
 \omega + \delta_1\kappa + a_1\kappa^3 - b_1\kappa^4 &= (\omega + \delta_2\kappa + a_2\kappa^3 - b_2\kappa^4)\chi, \\
 (e_1 + f_1\kappa)(c_1 + d_1\chi^2) + (\gamma_1 + \eta_1\kappa)\chi^2 - \lambda_1\kappa \\
 &= (e_2 + f_2\kappa)(c_2\chi^2 + d_2)\chi + (\gamma_2 + \eta_2\kappa)\chi - \lambda_2\kappa\chi^3.
 \end{aligned}
 \tag{18}$$

From (18), we have the wave number of the soliton as

$$\omega = \frac{\delta_1\kappa + a_1\kappa^3 - b_1\kappa^4 - (\delta_2\kappa + a_2\kappa^3 - b_2\kappa^4)\chi}{(\chi - 1)}.
 \tag{19}$$

Equation (10) can be rewritten as

$$b_1U_1'''' + \Pi_0U_1'' + \Pi_1U_1 + \Pi_3U_1^3 = 0,
 \tag{20}$$

where

$$\left. \begin{aligned}
 \Pi_0 &= 3a_1\kappa - 6b_1\kappa^2, \\
 \Pi_1 &= -[\omega + \delta_1\kappa + a_1\kappa^3 - b_1\kappa^4], \\
 \Pi_3 &= (e_1 + f_1\kappa)(c_1 + d_1\chi^2) + (\gamma_1 + \eta_1\kappa)\chi^2 - \lambda_1\kappa.
 \end{aligned} \right\}
 \tag{21}$$

Now, we will solve Eq. (20) using the following methods.

### 3. The Generalized Auxiliary Equation Method

According to this method, we assume that Eq. (20) has the formal solution:

$$U_1(\xi) = \sum_{l=0}^N \alpha_l f^l(\xi),
 \tag{22}$$

where  $f(\xi)$  satisfies the first-order auxiliary equation:

$$f'^2(\xi) = \sum_{m=0}^6 h_m f^m(\xi).
 \tag{23}$$

Here,  $\alpha_l (l = 0, 1, \dots, N)$  and  $h_m (m = 0, 1, \dots, 6)$  are constants to be determined such that  $\alpha_N \neq 0$  and  $h_6 \neq 0$ , where  $N$  is a positive integer. Balancing  $U_1''''$  and  $U_1^3$  in Eq. (20), one gets the balance number  $N = 4$ . From (22), one gets the formal solution of Eq. (20) as

$$U_1(\xi) = \alpha_0 + \alpha_1 f(\xi) + \alpha_2 f^2(\xi) + \alpha_3 f^3(\xi) + \alpha_4 f^4(\xi),
 \tag{24}$$

where  $\alpha_l (l = 0, 1, 2, 3, 4)$  are constants to be determined such that  $\alpha_4 \neq 0$ . Substituting (24) and (23) with  $h_0 = h_1 = h_3 = h_5 = 0$  into Eq. (20), collecting all the coefficients of  $[f(\xi)]^{m_1} [f(\xi)]^{m_2} (m_1 = 0, 1, \dots, 12, m_2 = 0, 1)$  and setting these coefficients to zero, we have a set of algebraic equations which can be solved using the

Maple to obtain the following results:

$$\left. \begin{aligned}
 &\Pi_3\alpha_4^3 + 1920b_1\alpha_4h_6^2 = 0, \\
 &3\Pi_3\alpha_3\alpha_4^2 + 945b_1\alpha_3h_6^2 = 0, \\
 &3\Pi_3\alpha_2\alpha_4^2 + 384b_1\alpha_2h_6^2 + 3\Pi_3\alpha_3^2\alpha_4 + 2688b_1\alpha_4h_6h_4 = 0, \\
 &105b_1\alpha_1h_6^2 + 1260b_1h_4\alpha_3h_6 + 6\Pi_3\alpha_2\alpha_3\alpha_4 + 3\Pi_3\alpha_1\alpha_4^2 + \Pi_3\alpha_3^3 = 0, \\
 &6\Pi_3\alpha_1\alpha_3\alpha_4 + 3\Pi_3\alpha_2\alpha_3^2 + 24\Pi_0\alpha_4h_6 + 3\Pi_3\alpha_0\alpha_4^2 + 1920b_1\alpha_4h_6h_2 \\
 &\quad + 840b_1\alpha_4h_4^2 + 3\Pi_3\alpha_2^2\alpha_4 + 480b_1h_4\alpha_2h_6 = 0, \\
 &870b_1h_6\alpha_3h_2 + 6\Pi_3\alpha_1\alpha_2\alpha_4 + 15\Pi_0\alpha_3h_6 + 3\Pi_3\alpha_1\alpha_3^2 + 360b_1\alpha_3h_4^2 \\
 &\quad + 6\Pi_3\alpha_0\alpha_3\alpha_4 + 120b_1h_4\alpha_1h_6 + 3\Pi_3\alpha_2^2\alpha_3 = 0, \\
 &320b_1h_6\alpha_2h_2 + 6\Pi_3\alpha_0\alpha_2\alpha_4 + 6\Pi_3\alpha_1\alpha_2\alpha_3 + 3\Pi_3\alpha_1^2\alpha_4 + 20\Pi_0\alpha_4h_4 \\
 &\quad + 8\Pi_0\alpha_2h_6 + \Pi_3\alpha_2^3 + 120b_1\alpha_2h_4^2 + 3\Pi_3\alpha_0\alpha_3^2 + 1040b_1h_4\alpha_4h_2 = 0, \\
 &24b_1\alpha_1h_4^2 + 78b_1h_6\alpha_1h_2 + 6\Pi_3\alpha_0\alpha_2\alpha_3 + 6\Pi_3\alpha_0\alpha_1\alpha_4 + 408b_1h_4\alpha_3h_2 \\
 &\quad + 3\Pi_3\alpha_1\alpha_2^2 + 12\Pi_0\alpha_3h_4 + 3\Pi_0\alpha_1h_6 + 3\Pi_3\alpha_1^2\alpha_3 = 0, \\
 &120b_1h_4\alpha_2h_2 + 6\Pi_3\alpha_0\alpha_1\alpha_3 + \Pi_1\alpha_4 + 6\Pi_0\alpha_2h_4 + 256b_1\alpha_4h_2^2 \\
 &\quad + 3\Pi_3\alpha_0^2\alpha_4 + 3\Pi_3\alpha_0\alpha_2^2 + 3\Pi_3\alpha_1^2\alpha_2 + 16\Pi_0\alpha_4h_2 = 0, \\
 &20b_1h_4\alpha_1h_2 + 9\Pi_0\alpha_3h_2 + 6\Pi_3\alpha_0\alpha_1\alpha_2 + 81b_1\alpha_3h_2^2 + 3\Pi_3\alpha_0^2\alpha_3 \\
 &\quad + \Pi_3\alpha_1^3 + 2\Pi_0\alpha_1h_4 + \Pi_1\alpha_3 = 0, \\
 &3\Pi_3\alpha_0^2\alpha_2 + 16b_1\alpha_2h_2^2 + 3\Pi_3\alpha_0\alpha_1^2 + 4\Pi_0\alpha_2h_2 + \Pi_1\alpha_2 = 0, \\
 &3\Pi_3\alpha_0^2\alpha_1 + b_1\alpha_1h_2^2 + \Pi_0\alpha_1h_2 + \Pi_1\alpha_1 = 0, \\
 &\Pi_1\alpha_0 + \Pi_3\alpha_0^3 = 0.
 \end{aligned} \right\} \quad (25)$$

On solving the above algebraic equations (25) by using the Maple, one gets the following two results:

Result 1:

$$\begin{aligned}
 \alpha_0 = 0, \quad \alpha_1 = 0, \quad \alpha_2 = 4\epsilon\sqrt{\frac{6\Pi_0h_6}{\Pi_3}}, \quad \alpha_3 = 0, \quad \alpha_4 = 8\epsilon h_6\sqrt{-\frac{30b_1}{\Pi_3}}, \\
 h_2 = -\frac{\Pi_0}{20b_1}, \quad h_4 = -\epsilon\sqrt{-\frac{\Pi_0h_6}{5b_1}}, \quad h_6 = h_6
 \end{aligned} \quad (26)$$

and

$$\Pi_1 = \frac{4\Pi_0^2}{25b_1}, \quad (27)$$

provided  $\Pi_0\Pi_3h_6 > 0$ ,  $\Pi_0b_1h_6 < 0$ ,  $b_1\Pi_3 < 0$  and  $\epsilon = \pm 1$ . It is well known that Eq. (23) has the following solutions: When  $h_0 = h_1 = h_3 = h_5 = 0$ , we have

$$f(\xi) = \left[ -\frac{h_2h_4\operatorname{sech}^2(\sqrt{h_2}\xi)}{h_4^2 - h_2h_6[1 + \epsilon \tanh(\sqrt{h_2}\xi)]^2} \right]^{\frac{1}{2}}, \quad (28)$$

$$f(\xi) = \left[ \frac{h_2 h_4 \operatorname{csch}^2(\sqrt{h_2} \xi)}{h_4^2 - h_2 h_6 [1 + \epsilon \coth(\sqrt{h_2} \xi)]^2} \right]^{\frac{1}{2}}, \quad (29)$$

provided  $h_2 > 0$ ,  $\epsilon = \pm 1$ ,

$$f(\xi) = \left[ -\frac{h_2 \operatorname{sech}^2(\sqrt{h_2} \xi)}{h_4 + 2\epsilon \sqrt{h_2 h_6} \tanh(\sqrt{h_2} \xi)} \right]^{\frac{1}{2}}, \quad (30)$$

$$f(\xi) = \left[ \frac{h_2 \operatorname{csch}^2(\sqrt{h_2} \xi)}{h_4 + 2\epsilon \sqrt{h_2 h_6} \coth(\sqrt{h_2} \xi)} \right]^{\frac{1}{2}}, \quad (31)$$

provided  $h_2 > 0$ ,  $h_6 > 0$  and  $\epsilon = \pm 1$ .

Substituting (26) along with (28)–(31) into (24), one gets the following soliton solutions of the system (1) and (2):

$$q(x, t) = \pm \frac{4}{5} \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \frac{\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2} \left[ 1 - \frac{2\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2} \right] e^{i(-\kappa x + \omega t + \theta_0)} \quad (32)$$

and

$$r(x, t) = \pm \frac{4}{5} \chi \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \frac{\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2} \left[ 1 - \frac{2\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2} \right] e^{i(-\kappa x + \omega t + \theta_0)}, \quad (33)$$

or

$$q(x, t) = \pm \frac{4}{5} \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \frac{\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2} \left[ 1 + \frac{2\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2} \right] e^{i(-\kappa x + \omega t + \theta_0)} \quad (34)$$

and

$$r(x, t) = \pm \frac{4}{5} \chi \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \frac{\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2} \left[1 + \frac{2\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{4 - \left[1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)\right]^2}\right] e^{i(-\kappa x + \omega t + \theta_0)}, \quad (35)$$

or

$$q(x, t) = \pm \frac{1}{2} \Pi_0 \sqrt{-\frac{6}{5b_1 \Pi_3}} \left[ \frac{\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] \left[ 2 - \frac{\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] e^{i(-\kappa x + \omega t + \theta_0)}, \quad (36)$$

and

$$r(x, t) = \pm \frac{1}{2} \chi \Pi_0 \sqrt{-\frac{6}{5b_1 \Pi_3}} \left[ \frac{\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] \left[ 2 - \frac{\operatorname{sech}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \tanh\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] e^{i(-\kappa x + \omega t + \theta_0)}, \quad (37)$$

or

$$q(x, t) = \pm \frac{1}{2} \Pi_0 \sqrt{-\frac{6}{5b_1 \Pi_3}} \left[ \frac{\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] \left[ 2 + \frac{\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] e^{i(-\kappa x + \omega t + \theta_0)}, \quad (38)$$

and

$$r(x, t) = \pm \frac{1}{2} \chi \Pi_0 \sqrt{-\frac{6}{5b_1 \Pi_3}} \left[ \frac{\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] \left[ 2 + \frac{\operatorname{csch}^2\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)}{1 + \epsilon \coth\left(\sqrt{-\frac{\Pi_0}{20b_1}}(x - vt)\right)} \right] e^{i(-\kappa x + \omega t + \theta_0)}, \quad (39)$$

provided  $b_1 \Pi_3 < 0$  and  $b_1 \Pi_0 < 0$ .



The solutions (32)–(39) exist under the constraint condition (27).

Result 2:

$$\alpha_0 = 0, \quad \alpha_1 = 0, \quad \alpha_2 = 0, \quad \alpha_3 = 0, \quad \alpha_4 = 8\epsilon h_6 \sqrt{-\frac{30b_1}{\Pi_3}}, \quad (40)$$

$$h_2 = -\frac{\Pi_0}{80b_1}, \quad h_4 = 0, \quad h_6 = h_6$$

and

$$\Pi_1 = \frac{4\Pi_0^2}{25b_1}, \quad (41)$$

provided  $b_1\Pi_3 < 0$  and  $\epsilon = \pm 1$ . It is well known that Eq. (23) has the following solutions:

When  $h_0 = h_1 = h_3 = h_5 = 0$ , we have

$$f(\xi) = \left[ \frac{2h_2}{\epsilon \sqrt{h_4^2 - 4h_2h_6} \cosh(2\sqrt{h_2}\xi) - h_4} \right]^{\frac{1}{2}}, \quad (42)$$

provided  $h_2 > 0, h_4^2 - 4h_2h_6 > 0$  and  $\epsilon = \pm 1$ ,

$$f(\xi) = \left[ \frac{2h_2}{\epsilon \sqrt{-(h_4^2 - 4h_2h_6)} \sinh(2\sqrt{h_2}\xi) - h_4} \right]^{\frac{1}{2}}, \quad (43)$$

provided  $h_2 > 0, h_4^2 - 4h_2h_6 < 0$  and  $\epsilon = \pm 1$ ,

$$f(\xi) = \left[ \frac{2h_2}{\epsilon \sqrt{h_4^2 - 4h_2h_6} \cos(2\sqrt{-h_2}\xi) - h_4} \right]^{\frac{1}{2}}, \quad (44)$$

$$f(\xi) = \left[ \frac{2h_2}{\epsilon \sqrt{h_4^2 - 4h_2h_6} \sin(2\sqrt{-h_2}\xi) - h_4} \right]^{\frac{1}{2}}, \quad (45)$$

provided  $h_2 < 0, h_4^2 - 4h_2h_6 > 0$  and  $\epsilon = \pm 1$ .

Substituting (40) along with (42)–(45) into (24), one gets the following solutions of the system (1) and (2):

(I) The bright soliton solutions:

$$q(x, t) = \pm \frac{1}{10} \Pi_0 \sqrt{-\frac{30}{b_1\Pi_3}} \operatorname{sech}^2 \left( \frac{1}{2} \sqrt{-\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)} \quad (46)$$

and

$$r(x, t) = \pm \frac{1}{10} \chi \Pi_0 \sqrt{-\frac{30}{b_1\Pi_3}} \operatorname{sech}^2 \left( \frac{1}{2} \sqrt{-\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)}, \quad (47)$$

provided  $b_1\Pi_3 < 0$  and  $b_1\Pi_0 < 0$ .

(II) The singular soliton solutions:

$$q(x, t) = \pm \frac{1}{10} \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \operatorname{csch}^2 \left( \frac{1}{2} \sqrt{-\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)} \quad (48)$$

and

$$r(x, t) = \pm \frac{1}{10} \chi \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \operatorname{csch}^2 \left( \frac{1}{2} \sqrt{-\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)}, \quad (49)$$

provided  $b_1 \Pi_3 < 0$  and  $b_1 \Pi_0 < 0$ .

(III) The periodic wave solutions:

$$q(x, t) = \pm \frac{1}{10} \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \operatorname{sec}^2 \left( \frac{1}{2} \sqrt{\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)} \quad (50)$$

and

$$r(x, t) = \pm \frac{1}{10} \chi \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \operatorname{sec}^2 \left( \frac{1}{2} \sqrt{\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)}, \quad (51)$$

or

$$q(x, t) = \pm \frac{1}{10} \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \operatorname{csc}^2 \left( \frac{1}{2} \sqrt{\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)} \quad (52)$$

and

$$r(x, t) = \pm \frac{1}{10} \chi \Pi_0 \sqrt{-\frac{30}{b_1 \Pi_3}} \operatorname{csc}^2 \left( \frac{1}{2} \sqrt{\frac{\Pi_0}{5b_1}} (x - vt) \right) e^{i(-\kappa x + \omega t + \theta_0)}, \quad (53)$$

provided  $b_1 \Pi_3 < 0$  and  $b_1 \Pi_0 > 0$ .

The solutions (46)–(53) exist under the constraint condition (41).

#### 4. An Addendum to Kudryashov’s Method

According to this method, we assume that Eq. (20) has the formal solution

$$U_1(\xi) = \sum_{l=0}^N B_l R^l(\xi), \quad (54)$$

where  $B_l (l = 0, 1, 2, \dots, N)$  are constants to be determined, such that  $B_N \neq 0$ , while  $R(\xi)$  satisfies the first-order auxiliary ordinary differential equation (ODE):

$$R^2(\xi) = R^2(\xi)[1 - \Gamma R^{2p}(\xi)] \ln^2 K, \quad 0 < K \neq 1, \quad (55)$$

where  $\Gamma$  is a constant. It is easy to show that Eq. (55) has the following solution:

$$R(\xi) = \left[ \frac{4A}{4A^2 \exp_K(p\xi) + \Gamma \exp_K(-p\xi)} \right]^{\frac{1}{p}}, \tag{56}$$

where  $A$  is a nonzero constant,  $p$  is a positive integer and  $\exp_K(p\xi) = K^{p\xi}$ . We determine the positive integer  $N$  in (54) by using the homogeneous balance method as follows.

If  $D(U_1) = N, D(U_1') = N + p, D(U_1'') = N + 2p$ , then we have  $D[U_1^r U_1^{(s)}] = N(r + 1) + ps$ .

Now, by balancing  $U_1''''$  with  $U_1^3$  in Eq. (20), one gets

$$N + 4p = 3N \Rightarrow N = 2p. \tag{57}$$

Next, we will discuss the following cases.

**Case 1.** If we choose  $p = 1$ , then  $N = 2$ . Thus, from (54) we deduce that Eq. (20) has the formal solution:

$$U_1(\xi) = B_0 + B_1 R(\xi) + B_2 R^2(\xi), \tag{58}$$

where  $B_0, B_1$  and  $B_2$  are constants to be determined, such that  $B_2 \neq 0$  while  $R(\xi)$  satisfies the first-order auxiliary ODE:

$$R'^2(\xi) = R^2(\xi)[1 - \Gamma R^2(\xi)] \ln^2 K, \quad 0 < K \neq 1, \tag{59}$$

where  $\Gamma$  is a constant. Substituting (58) along with (59) into Eq. (20), collecting all the coefficients of each power of  $[R(\xi)]^{m_1} [R'(\xi)]^{m_2}$  ( $m_1 = 0, 1, \dots, 6, m_2 = 0, 1$ ) and setting each of these coefficients to zero, we have a system of algebraic equations which can be solved by using the Maple to get the results:

$$B_0 = 0, \quad B_1 = 0, \quad B_2 = \pm 2\Gamma \sqrt{-\frac{30b_1}{\Pi_3} \ln^2 K} \tag{60}$$

and

$$\Pi_0 = -20b_1 \ln^2 K, \quad \Pi_1 = 64b_1 \ln^4 K, \tag{61}$$

provided  $b_1 \Pi_3 < 0$ . Substituting (60) along with (56) into Eq. (58), one gets the solutions of the system (1) and (2) in the forms:

$$q(x, t) = \pm 2\Gamma \sqrt{-\frac{30b_1}{\Pi_3}} \left[ \frac{4A \ln K}{4A^2 K^{(x-ct)} + \Gamma K^{-(x-ct)}} \right]^2 e^{i(-\kappa x + \omega t + \theta_0)} \tag{62}$$

and

$$r(x, t) = \pm 2\chi \Gamma \sqrt{-\frac{30b_1}{\Pi_3}} \left[ \frac{4A \ln K}{4A^2 K^{(x-ct)} + \Gamma K^{-(x-ct)}} \right]^2 e^{i(-\kappa x + \omega t + \theta_0)}. \tag{63}$$

In particular, if we set  $\Gamma = 4A^2$  in (62) and (63), then the system (1) and (2) has the bright soliton solutions as

$$q(x, t) = \pm 2\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{sech}^2[(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)} \quad (64)$$

and

$$r(x, t) = \pm 2\chi\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{sech}^2[(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)}, \quad (65)$$

while, if  $\Gamma = -4A^2$ , one gets the singular soliton solution of the system (1) and (2) as

$$q(x, t) = \pm 2\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{csch}^2[(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)} \quad (66)$$

and

$$r(x, t) = \pm 2\chi\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{csch}^2[(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)}. \quad (67)$$

The solutions (62)–(67) exist under the conditions (61).

**Case 2.** If we choose  $p = 2$ , then  $N = 4$ . Thus, we deduce from (54) that Eq. (20) has the formal solution:

$$U_1(\xi) = B_0 + B_1R(\xi) + B_2R^2(\xi) + B_3R^3(\xi) + B_4R^4(\xi), \quad (68)$$

where  $B_0, B_1, B_2, B_3$  and  $B_4$  are constants to be determined, such that  $B_4 \neq 0$  and the function  $R(\xi)$  satisfies the first-order auxiliary ODE

$$R'(\xi) = R^2(\xi)[1 - \Gamma R^4(\xi)]\ln^2 K, \quad 0 < K \neq 1, \quad (69)$$

where  $\Gamma$  is a constant. Substituting (68) along with (69) into Eq. (20), collecting all the coefficients of each power of  $[R(\xi)]^{m_1}[R'(\xi)]^{m_2}$  ( $m_1 = 0, 1, 2, \dots, 12, m_2 = 0, 1$ ) and setting each of these coefficients to zero, we have a system of algebraic equations which can be solved by using the Maple to get the results:

$$B_0 = 0, \quad B_1 = 0, \quad B_2 = 0, \quad B_3 = 0, \quad B_4 = \pm 8\Gamma\sqrt{-\frac{30b_1}{\Pi_3}}\ln^2 K \quad (70)$$

and

$$\Pi_0 = -80b_1\ln^2 K, \quad \Pi_1 = 1024b_1\ln^4 K, \quad (71)$$

provided  $b_1\Pi_3 < 0$ . Substituting (70) along with (56) into Eq. (68), one gets the solutions of the system (1) and (2) in the forms:

$$q(x, t) = \pm 8\Gamma\sqrt{-\frac{30b_1}{\Pi_3}}\left[\frac{4A \ln K}{4A^2 K^{2(x-ct)} + \Gamma K^{-2(x-ct)}}\right]^2 e^{i(-\kappa x + \omega t + \theta_0)} \quad (72)$$

and

$$r(x, t) = \pm 8\chi\Gamma\sqrt{-\frac{30b_1}{\Pi_3}}\left[\frac{4A \ln K}{4A^2K^{2(x-ct)} + \Gamma K^{-2(x-ct)}}\right]^2 e^{i(-\kappa x + \omega t + \theta_0)}. \quad (73)$$

In particular, if we set  $\Gamma = 4A^2$  in (72) and (73), then we have the bright soliton solution of the system (1) and (2) as

$$q(x, t) = \pm 8\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{sech}^2[2(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)} \quad (74)$$

and

$$r(x, t) = \pm 8\chi\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{sech}^2[2(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)}, \quad (75)$$

while, if  $\Gamma = -4A^2$ , one gets the singular soliton solution of the system (1) and (2) as

$$q(x, t) = \pm 8\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{csch}^2[2(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)} \quad (76)$$

and

$$r(x, t) = \pm 8\chi\sqrt{-\frac{30b_1}{\Pi_3}}(\ln^2 K)\operatorname{csch}^2[2(x - ct) \ln K]e^{i(-\kappa x + \omega t + \theta_0)}. \quad (77)$$

The solutions (72)–(77) exist under the conditions (71).

Similarly, we can find many other solutions by choosing other values for  $p$  and  $N$ .

### 5. Conservation Laws

Following some detailed calculations, we get that system (1) and (2) admits three conservation laws corresponding to “power”, “linear momentum” and “Hamiltonian” for particular cases of the parameters; the conserved densities are, respectively, given as follows:

(i) If  $\eta_1 = \eta_2$  and  $d_1 = d_2 = 0$ , we get the conserved “power density”

$$T_c^t = \frac{1}{2}(|q|^2 + |r|^2). \quad (78)$$

(ii) If  $\eta_1 = \eta_2$ ,  $d_1 = d_2 = 0$ ,  $\lambda_1 = -\mu_2$ ,  $\lambda_2 = -\mu_2$  and  $\gamma_1 = \gamma_2$ , the conserved “linear momentum density” is

$$T_m^t = \frac{1}{2}(\mathcal{I}(q^*q_x) + \mathcal{I}(r^*r_x)). \quad (79)$$

(iii) If, in addition to the conditions in (ii),  $\eta_1 = \eta_2 = 0$ , the conserved “Hamiltonian density” is

$$\begin{aligned}
 T_e^t = & -a_1 \mathcal{I}(q^* q_{xxx}) - a_2 \mathcal{I}(r^* r_{xxx}) + b_1 \mathcal{R}(qq_{xxx}^*) + b_2 \mathcal{R}(rr_{xxx}^*) \\
 & + \frac{1}{4} c_1 |q|^2 (e_1 |q|^2 - f_1 \mathcal{I}(q^* q_x)) + \frac{1}{4} c_2 |r|^2 (e_2 |r|^2 - f_2 \mathcal{I}(r^* r_x)) \\
 & + \frac{1}{2} \delta_1 \mathcal{I}(q^* q_x) + \frac{1}{2} \delta_2 \mathcal{I}(r^* r_x) - \frac{1}{4} \mu_1 |q|^2 \mathcal{I}(q^* q_x) - \frac{1}{4} \mu_2 |r|^2 \mathcal{I}(r^* r_x) \\
 & + \frac{1}{2} \gamma_2 |q|^2 |r|^2.
 \end{aligned} \tag{80}$$

The two-component bright soliton solutions recovered in this paper are structured as

$$q(x, t) = A_1 \operatorname{sech}^2[B(x - vt)] e^{i(-\kappa x + \omega t + \theta_0)}, \tag{81}$$

$$r(x, t) = A_2 \operatorname{sech}^2[B(x - vt)] e^{i(-\kappa x + \omega t + \theta_0)}. \tag{82}$$

Here,  $A_j$  ( $j = 1, 2$ ) are the amplitudes of the two solitons while  $B$  represents their inverse widths. Therefore, the conserved quantities are

$$P = \frac{1}{2} \int_{-\infty}^{\infty} (|q|^2 + |r|^2) dx = \frac{2}{3B} (|A_1|^2 + |A_2|^2), \tag{83}$$

$$M = \frac{1}{4i} \int_{-\infty}^{\infty} \{(q^* q_x - qq_x^*) + (r^* r_x - rr_x^*)\} dx = -\frac{2\kappa}{3B} (|A_1|^2 + |A_2|^2) \tag{84}$$

and

$$\begin{aligned}
 H = & \int_{-\infty}^{\infty} \left[ -\frac{1}{2i} \{a_1 (q^* q_{xxx} - qq_{xxx}^*) + a_2 (r^* r_{xxx} - rr_{xxx}^*)\} \right. \\
 & + \frac{1}{2} \{b_1 (q^* q_{xxxx} + qq_{xxxx}^*) + b_2 (r^* r_{xxxx} + rr_{xxxx}^*)\} \\
 & - \frac{1}{8i} \{(c_1 f_1 + \mu_1) |q|^2 (q^* q_x - qq_x^*) + (c_2 f_2 + \mu_2) |r|^2 (r^* r_x - rr_x^*)\} \\
 & \left. + \frac{1}{4i} \{\delta_1 (q^* q_x - qq_x^*) + \delta_2 (r^* r_x - rr_x^*)\} + \frac{\gamma_2}{2} |q|^2 |r|^2 \right] dx \\
 = & -\frac{4\kappa}{15} (a_1 A_1^2 + a_2 A_2^2) (5\kappa^2 + 12B^2) + \frac{64}{21} B^5 (b_1 A_1^2 + b_2 A_2^2) \\
 & + \frac{224}{35} \kappa^2 B (b_1 A_1^4 + b_2 A_2^4) + \frac{4\kappa^4}{3B} (b_1 A_1^5 + b_2 A_2^5) \\
 & + \frac{8\kappa}{35B} \{(c_1 f_1 + \mu_1) A_1^4 + (c_2 f_2 + \mu_2) A_2^4\} - \frac{2\kappa}{3B} (\delta_1 A_1^2 + \delta_2 A_2^2) + \frac{16\gamma_2 A_1^2 A_2^2}{35B}.
 \end{aligned} \tag{85}$$

These conserved quantities are power ( $P$ ), linear momentum ( $M$ ) and Hamiltonian ( $H$ ), respectively.

## 6. Conclusions

Today's work is a display of polarized soliton solutions and their conservation laws with FLE in birefringent fibers. Two integration algorithms have made this retrieval of soliton solutions possible while the approach of multipliers has successfully recovered the conservation laws. The conserved quantities are finally computed and exhibited by utilizing the soliton solutions that are recovered in the paper. These recovered and enumerated results serve as a gateway to extend the study further along in this direction. An immediate example would be to extend the model to address DWDM topology with FLE, studying the cascaded system and Thirring solitons. More results, aligned with recently reported works, will appear over time sequentially.<sup>28–39</sup> Consequently, the aspects of ghost pulses and suppression of intra-channel collision of solitons, stochastic perturbation of solitons are all on the table. These projects will be taken up and their results would be disseminated.

## Disclosure

The authors also declare that there is no conflict of interest.

## References

1. K. S. Al-Ghafri, E. V. Krishnan and A. Biswas, Chirped optical soliton perturbation of Fokas–Lenells equation with full nonlinearity, *Adv. Differ. Equ.* **2020** (2020) 191.
2. S. Arshed, A. Biswas, Q. Zhou, S. P. Moshokoa and M. Belic, Optical solitons with polarization–mode dispersion for coupled Fokas–Lenells equation with two forms of integration architecture, *Opt. Quantum Electron.* **50** (2018) 304.
3. A. Bansal, A. H. Kara, A. Biswas, S. P. Moshokoa and M. Belic, Optical soliton perturbation, group invariants and conservation laws of perturbed Fokas–Lenells equation, *Chaos Solitons Fractals* **114** (2018) 275–280.
4. A. Bansal, A. H. Kara, A. Biswas, S. Khan, Q. Zhou and S. P. Moshokoa, Optical solitons and conservation laws with polarization–mode dispersion for coupled Fokas–Lenells equation using group invariance, *Chaos Solitons Fractals* **120** (2019) 245–249.
5. A. Biswas, Chirp-free bright optical soliton perturbation with Fokas–Lenells equation by traveling wave hypothesis and semi-inverse variational principle, *Optik* **170** (2018) 431–435.
6. A. Biswas, Y. Yıldırım, E. Yaşar, Q. Zhou, S. P. Moshokoa and M. Belic, Optical soliton solutions to Fokas–Lenells equation using some different methods, *Optik* **173** (2018) 21–31.
7. A. Biswas, M. Ekici, A. Sonmezoglu and R. T. Alqahtani, Optical soliton perturbation with full nonlinearity for Fokas–Lenells equation, *Optik* **165** (2018) 29–34.
8. A. Biswas, Y. Yıldırım, E. Yaşar, Q. Zhou, M. F. Mahmood, S. P. Moshokoa and M. Belic, Optical solitons with differential group delay for coupled Fokas–Lenells equation using two integration schemes, *Optik* **165** (2018) 74–86.
9. A. Biswas, M. Ekici, A. Sonmezoglu and R. T. Alqahtani, Optical solitons with differential group delay for coupled Fokas–Lenells equation by extended trial function scheme, *Optik* **165** (2018) 102–110.

10. A. Biswas, H. Rezazadeh, M. Mirzazadeh, M. Eslami, M. Ekici, Q. Zhou, S. P. Moshokoa and M. Belic, Optical soliton perturbation with Fokas–Lenells equation using three exotic and efficient integration schemes, *Optik* **165** (2018) 288–294.
11. A. Biswas and S. Arshed, Application of semi-inverse variational principle to cubic–quartic optical solitons having Kerr and power law non-linearity, *Optik* **172** (2018) 847–50.
12. O. Gonzalez-Gaxiola, A. Biswas, F. Mallawi and M. R. Belic, Cubic–quartic bright optical solitons with improved Adomian decomposition method, *J. Adv. Res.* **21** (2020) 161–167.
13. K. Hosseini, M. Mirzazadeh, J. Vahidi and R. Asghari, Optical wave structures to the Fokas—Lenells equation, *Optik* **207** (2020) 164450.
14. A. J. M. Jawad, A. Biswas, Q. Zhou, S. P. Moshokoa and M. Belic, Optical soliton perturbation of Fokas–Lenells equation with two integration schemes, *Optik* **165** (2018) 111–116.
15. E. V. Krishnan, A. Biswas, Q. Zhou and M. Alifras, Optical soliton perturbation with Fokas–Lenells equation by mapping methods, *Optik* **178** (2019) 104–110.
16. N. A. Kudryashov, First integrals and general solution of the Fokas–Lenells equation, *Optik* **195** (2019) 163135.
17. N. A. Kudryashov, A re-visitation of the results on Fokas–Lenells equation by Mahak and Akram, *Optik* **209** (2020) 164522.
18. Y. Xiao, E. Fan and J. Xu, The Fokas–Lenells equation on the finite interval, *Acta Math. Sci.* **37** (2017) 852–876.
19. H. Triki and A. M. Wazwaz, Combined optical solitary waves of the Fokas–Lenells equation, *Waves Random Complex Media* **27** (2017) 587–593.
20. H. Triki and A. M. Wazwaz, New types of chirped soliton solutions for the Fokas–Lenells equation, *Int. J. Numer. Methods Heat Fluid Flow* **27** (2017) 1596–1601.
21. Y. Yildirim, A. Biswas, A. J. M. Jawad, M. Ekici, Q. Zhou, S. Khan, A. K. Alzahrani and M. R. Belic, Cubic–quartic optical solitons in birefringent fibers with four forms of nonlinear refractive index by exp-function expansion, *Results Phys.* **16** (2020) 102913.
22. E. M. E. Zayed, M. El-Horbaty and M. E. M. Alngar, Cubic–quartic optical soliton perturbation having four laws nonlinearity with a prolific integration algorithm, *Optik* **220** (2020) 165121.
23. E. M. E. Zayed, M. E. M. Alngar, M. M. El-Horbaty, A. Biswas, M. Ekici, Q. Zhou and M. R. Belic, Highly dispersive optical solitons in the nonlinear Schrödinger’s equation having polynomial law of the refractive index change, *Indian J. Phys.* **95**(1) (2021) 109–119.
24. E. M. E. Zayed, R. M. A. Shohib, M. E. M. Alngar, A. Biswas, M. Ekici, S. Khan, A. K. Alzahrani and M. R. Belic, Optical solitons and conservation laws associated with Kudryashov’s sextic power-law nonlinearity of refractive index, *Uk. J. Phys. Opt.* **22**(1) (2021) 38–49.
25. Q. Zhou and A. Biswas, Optical solitons in parity–time symmetric mixed linear and nonlinear lattice with non-Kerr law nonlinearity, *Superlattices Microstruct.* **109** (2017) 588–598.
26. Q. Zhou, A. H. Arnous, S. P. Moshokoa, M. Z. Ullah, M. Ekici, M. Mirzazadeh, A. S. Alshomrani, A. Biswas and M. Belic, Optical solitons in DWDM system with modified simple equation method, *J. Optoelectron. Adv. Mater.* **20**(1–2) (2018) 46–51.
27. Q. Zhou, M. Mirzazadeh, M. Ekici, S. P. Moshokoa, A. Biswas, M. Z. Ullah and M. Belic, Dark and singular optical solitons with spatio–temporal dispersion by Kudryashov’s method, *J. Optoelectron. Adv. Mater.* **20**(3–4) (2018) 126–132.
28. N. A. Kudryashov, Highly dispersive solitary wave solutions of perturbed nonlinear Schrödinger equations, *Appl. Math. Comput.* **371** (2020) 124972.



29. N. A. Kudryashov, Method for finding highly dispersive optical solitons of nonlinear differential equations, *Optik* **206** (2020) 163550.
30. N. A. Kudryashov, Almost general solution of the reduced higher-order nonlinear Schrödinger equation, *Optik* **230** (2021) 166347.
31. N. A. Kudryashov, Solitary waves of the non-local Schrödinger equation with arbitrary refractive index, *Optik* **231** (2021) 166443.
32. N. A. Kudryashov, Solitary waves of the generalized Sasa–Satsuma equation with arbitrary refractive index, *Optik* **232** (2021) 166540.
33. N. A. Kudryashov, Solitary wave solutions of hierarchy with non-local nonlinearity, *Appl. Math. Lett.* **103** (2020) 106155.
34. J. Dan, S. Sain, A. Ghose-Choudhury and S. Garai, Application of the Kudryashov function for finding solitary wave solutions of NLS type differential equations, *Optik* **224** (2020) 165519.
35. S. Sain, A. Ghose-Choudhury and S. Garai, Solitary wave solutions for the KdV-type equations in plasma: A new approach with the Kudryashov function, *Eur. Phys. J. Plus* **136**(2) (2021) 226.
36. Y. Yildirim, A. Biswas, A. H. Kara, M. Ekici, S. Khan and M. R. Belic, Optical soliton perturbation and conservation law with Kudryashov’s refractive index having quadrupled power-law and dual form of generalized nonlocal nonlinearity, *Semicond. Phys. Quantum Electron. Optoelectron.* **24**(1) (2021) 64–70.
37. A. Biswas, Optical soliton cooling with polynomial law of nonlinear refractive index, *J. Opt.* **49**(4) (2020) 580–583.
38. E. M. E. Zayed, M. E. M. Alngar, A. Biswas, A. H. Kara, L. Moraru, M. Ekici, A. K. Alzahrani and M. R. Belic, Solitons and conservation laws in magneto-optic waveguides with triple-power law nonlinearity, *J. Opt.* **49**(4) (2020) 584–590.
39. E. M. E. Zayed, A. G. Al-Nowehy, M. E. M. Alngar, A. Biswas, M. Asma, M. Ekici, A. K. Alzahrani and M. R. Belic, Highly dispersive optical solitons in birefringent fibers with four nonlinear forms using Kudryashov’s approach, *J. Opt.* **50**(1) (2021) 120–131.